

Ca

3

The variation function and the absorption curve in the optical dissociation of thallium iodide. J. M. FRANK. *Physik. Z. Supplementum* 2, 310 (1932). The efficiency of excitation of Tl lines by optical disson. of TlI as a function of the exciting light was measured. This excitation function was compared with the curve of absorption coeff. of TlI in the range 186 to 226.5 m μ . It is shown that 3 processes must be assumed, to explain the absorption curve in this range, i. e., the disson. of TlI into Tl in the 295% condition + 1 into Tl' + 1- and still another process in the range of 180 m μ . C. F. P. JEFFREYS. Zhur. Fiz.

ALL INFORMATION CONTAINED HEREIN IS UNCLASSIFIED

APPROVED FOR RELEASE: 06/13/2000

CIA-RDP86-00513R000413530013-1"

1ST AND 2ND COLUMNS																										3RD AND 4TH COLUMNS																									
PROCESSES AND PROPERTIES INDEX																																																			
<p><i>OK</i></p> <p><i>Do not answer</i></p> <p>Oxidation of mercury vapors under the influence of ultra-violet light. I. M. Frank. <i>Compt. rend. acad. sci. U. R. S. S. [N. S.]</i>, 1943, 146-7 (in English 147-8); cf. Dickinson and Sherrill, <i>C. A.</i> 20, 1954; Noyes, <i>C. A.</i> 22, 2717; 23, 3165; Leipunskii and Segulin, <i>C. A.</i> 23, 1054, 5190.—In the oxidation of Hg vapors with O under the influence of ultra-violet, e. g., exciting the Hg atoms with the line 2537 Å., the reaction may follow the following course: $\text{Hg}^* + \text{O}_2 \rightarrow \text{Hg} + \text{O}_2^*$; $\text{O}_2^* + \text{O}_2 \rightarrow \text{O}_3 + \text{O}$; $\text{O}_3 + \text{Hg} \rightarrow \text{Hg}(\text{O}) + \text{O}_2$, where O_2^* is an excited mol. in the state $^3\Sigma$ or $^1\Sigma$. The expl. procedure carried out under sub-atm. pressure measured manometrically confirmed this assumption. The removal of short-wave light from the exciting rays has no effect on the reaction, but a decrease in the exciting rays has an adverse effect. The most effective lines are those of Al at 1880, 1815 and 1801 Å., which induce the formation of O_3 and Hg is not needed under the above conditions provided that some O_2 is already present.</p> <p>A. A. Boettlinck</p>																																																			
<p>ASS. S. A. METALLURGICAL LITERATURE CLASSIFICATION</p>																																																			

CO

PROCESSED AND PROPERTIES

Absorption of light near the mercury resonance line
 1850 Å. I. M. Frank. *Physik. Z. Sowjetunion* 6, 517-55
 (1933) (in German).—The absorption of the Al line at
 1454 Å 2 l. by Hg is proportional to the thickness of Hg
 vapor, and is 0.53 at 200° for a layer of Hg 10 mm. thick.
 Absorption decreased with increase in temp. The ab-
 sorption is due to scattering of the light by the Hg atom
 because of the close proximity to the 1449.6 Å. line of
 Hg.
 Gerald M. Petty

12

ASB-SLA METALLURGICAL LITERATURE CLASSIFICATION

1930-1939

1940-1949

1950-1959

1960-1969

1970-1979

1980-1989

1990-1999

2000-2009

2010-2019

2020-2029

2030-2039

2040-2049

2050-2059

2060-2069

2070-2079

2080-2089

2090-2099

2100-2109

2110-2119

2120-2129

2130-2139

2140-2149

2150-2159

2160-2169

2170-2179

2180-2189

2190-2199

2200-2209

2210-2219

2220-2229

2230-2239

2240-2249

2250-2259

2260-2269

2270-2279

2280-2289

2290-2299

2300-2309

2310-2319

2320-2329

2330-2339

2340-2349

2350-2359

2360-2369

2370-2379

2380-2389

2390-2399

2400-2409

2410-2419

2420-2429

2430-2439

2440-2449

2450-2459

2460-2469

2470-2479

2480-2489

2490-2499

2500-2509

2510-2519

2520-2529

2530-2539

2540-2549

2550-2559

2560-2569

2570-2579

2580-2589

2590-2599

2600-2609

2610-2619

2620-2629

2630-2639

2640-2649

2650-2659

2660-2669

2670-2679

2680-2689

2690-2699

2700-2709

2710-2719

2720-2729

2730-2739

2740-2749

2750-2759

2760-2769

2770-2779

2780-2789

2790-2799

2800-2809

2810-2819

2820-2829

2830-2839

2840-2849

2850-2859

2860-2869

2870-2879

2880-2889

2890-2899

2900-2909

2910-2919

2920-2929

2930-2939

2940-2949

2950-2959

2960-2969

2970-2979

2980-2989

2990-2999

3000-3009

3010-3019

3020-3029

3030-3039

3040-3049

3050-3059

3060-3069

3070-3079

3080-3089

3090-3099

3100-3109

3110-3119

3120-3129

3130-3139

3140-3149

3150-3159

3160-3169

3170-3179

3180-3189

3190-3199

3200-3209

3210-3219

3220-3229

3230-3239

3240-3249

3250-3259

3260-3269

3270-3279

3280-3289

3290-3299

3300-3309

3310-3319

3320-3329

3330-3339

3340-3349

3350-3359

3360-3369

3370-3379

3380-3389

3390-3399

3400-3409

3410-3419

3420-3429

3430-3439

3440-3449

3450-3459

3460-3469

3470-3479

3480-3489

3490-3499

3500-3509

3510-3519

3520-3529

3530-3539

3540-3549

3550-3559

3560-3569

3570-3579

3580-3589

3590-3599

3600-3609

3610-3619

3620-3629

3630-3639

3640-3649

3650-3659

3660-3669

3670-3679

3680-3689

3690-3699

3700-3709

3710-3719

3720-3729

3730-3739

3740-3749

3750-3759

3760-3769

3770-3779

3780-3789

3790-3799

3800-3809

3810-3819

3820-3829

3830-3839

3840-3849

3850-3859

3860-3869

3870-3879

3880-3889

3890-3899

3900-3909

3910-3919

3920-3929

3930-3939

3940-3949

3950-3959

3960-3969

3970-3979

3980-3989

3990-3999

4000-4009

4010-4019

4020-4029

4030-4039

4040-4049

4050-4059

4060-4069

4070-4079

4080-4089

4090-4099

4100-4109

4110-4119

4120-4129

4130-4139

4140-4149

4150-4159

4160-4169

4170-4179

4180-4189

4190-4199

4200-4209

4210-4219

4220-4229

4230-4239

4240-4249

4250-4259

4260-4269

4270-4279

4280-4289

4290-4299

4300-4309

4310-4319

4320-4329

4330-4339

4340-4349

4350-4359

4360-4369

4370-4379

4380-4389

4390-4399

4400-4409

4410-4419

4420-4429

4430-4439

4440-4449

4450-4459

4460-4469

4470-4479

4480-4489

4490-4499

4500-4509

4510-4519

4520-4529

4530-4539

4540-4549

4550-4559

4560-4569

4570-4579

4580-4589

4590-4599

4600-4609

4610-4619

4620-4629

4630-4639

4640-4649

4650-4659

4660-4669

4670-4679

4680-4689

4690-4699

4700-4709

4710-4719

4720-4729

4730-4739

4740-4749

4750-4759

4760-4769

4770-4779

4780-4789

4790-4799

4800-4809

4810-4819

4820-4829

4830-4839

4840-4849

4850-4859

4860-4869

4870-4879

4880-4889

4890-4899

4900-4909

4910-4919

4920-4929

4930-4939

4940-4949

4950-4959

4960-4969

4970-4979

4980-4989

4990-4999

5000-5009

5010-5019

5020-5029

5030-5039

5040-5049

5050-5059

5060-5069

5070-5079

5080-5089

5090-5099

5100-5109

5110-5119

5120-5129

5130-5139

5140-5149

5150-5159

5160-5169

5170-5179

5180-5189

5190-5199

5200-5209

5210-5219

5220-5229

5230-5239

5240-5249

5250-5259

5260-5269

5270-5279

5280-5289

5290-5299

5300-5309

5310-5319

5320-5329

5330-5339

5340-5349

5350-5359

5360-5369

5370-5379

5380-5389

5390-5399

5400-5409

5410-5419

5420-5429

5430-5439

5440-5449

5450-5459

5460-5469

5470-5479

5480-5489

5490-5499

5500-5509

5510-5519

5520-5529

5530-5539

5540-5549

5550-5559

5560-5569

5570-5579

5580-5589

5590-5599

5600-5609

5610-5619

5620-5629

5630-5639

5640-5649

5650-5659

5660-5669

5670-5679

5680-5689

5690-5699

5700-5709

5710-5719

5720-5729

5730-5739

5740-5749

5750-5759

5760-5769

5770-5779

5780-5789

5790-5799

5800-5809

5810-5819

5820-5829

5830-5839

5840-5849

5850-5859

5860-5869

5870-5879

5880-5889

5890-5899

5900-5909

5910-5919

5920-5929

5930-5939

5940-5949

5950-5959

5960-5969

5970-5979

5980-5989

5990-5999

6000-6009

6010-6019

6020-6029

6030-6039

6040-6049

6050-6059

6060-6069

6070-6079

6080-6089

6090-6099

6100-6109

6110-6119

6120-6129

6130-6139

6140-6149

6150-6159

6160-6169

6170-6179

6180-6189

6190-6199

6200-6209

6210-6219

6220-6229

6230-6239

6240-6249

6250-6259

6260-6269

6270-6279

6280-6289

6290-6299

6300-6309

6310-6319

6320-6329

6330-6339

6340-6349

6350-6359

6360-6369

6370-6379

6380-6389

6390-6399

6400-6409

6410-6419

6420-6429

6430-6439

6440-6449

6450-6459

6460-6469

6470-6479

6480-6489

6490-6499

6500-6509

6510-6519

6520-6529

6530-6539

6540-6549

6550-6559

6560-6569

6570-6579

6580-6589

6590-6599

6600-6609

6610-6619

6620-6629

6630-6639

6640-6649

6650-6659

6660-6669

6670-6679

6680-6689

6690-6699

6700-6709

6710-6719

6720-6729

6730-6739

6740-6749

6750-6759

6760-6769

6770-6779

6780-6789

6790-6799

6800-6809

6810-6819

6820-6829

6830-6839

6840-6849

6850-6859

6860-6869

6870-6879

6880-6889

6890-6899

6900-6909

6910-6919

6920-6929

6930-6939

6940-6949

6950-6959

6960-6969

6970-6979

6980-6989

6990-6999

7000-7009

7010-7019

7020-7029

7030-7039

7040-7049

7050-7059

7060-7069

7070-7079

7080-7089

7090-7099

7100-7109

7110-7119

7120-7129

7130-7139

7140-7149

7150-7159

7160-7169

7170-7179

7180-7189

7190-7199

7200-7209

7210-7219

7220-7229

7230-7239

7240-7249

7250-7259

7260-7269

7270-7279

7280-7289

7290-7299

7300-7309

7310-7319

7320-7329

7330-7339

7340-7349

7350-7359

7360-7369

7370-7379

7380-7389

7390-7399

7400-7409

7410-7419

7420-7429

7430-7439

7440-7449

7450-7459

7460-7469

7470-7479

7480-7489

7490-7499

7500-7509

7510-7519

7520-7529

7530-7539

7540-7549

7550-7559

7560-7569

7570-7579

7580-7589

7590-7599

7600-7609

7610-7619

7620-7629

7630-7639

7640-7649

7650-7659

7660-7669

7670-7679

7680-7689

7690-7699

7700-7709

7710-7719

7720-7729

7730-7739

7740-7749

7750-7759

7760-7769

7770-7779

7780-7789

7790-7799

7800-7809

7810-7819

7820-7829

7830-7839

7840-7849

7850-7859

7860-7869

7870-7879

7880-7889

7890-7899

7900-7909

7910-7919

7920-7929

7930-7939

7940-7949

7950-7959

7960-7969

7970-7979

7980-7989

7990-7999

8000-8009

8010-8019

8020-8029

8030-8039

8040-8049

8050-8059

8060-8069

8070-8079

8080-8089

8090-8099

8100-8109

8110-8119

8120-8129

8130-8139

8140-8149

8150-8159

8160-8169

8170-8179

8180-8189

8190-8199

8200-8209

8210-8219

8220-8229

8230-8239

8240-8249

8250-8259

8260-8269

8270-8279

8280-8289

8290-8299

8300-8309

8310-8319

8320-8329

8330-8339

8340-8349

8350-8359

8360-8369

8370-8379

8380-8389

8390-8399

8400-8409

8410-8419

8420-8429

8430-8439

8440-8449

8450-8459

8460-8469

8470-8479

8480-8489

8490-8499

8500-8509

8510-8519

8520-8529

8530-8539

8540-8549

8550-8559

8560-8569

8570-8579

8580-8589

8590-8599

8600-8609

8610-8619

8620-8629

8630-8639

8640-8649

8650-8659

8660-8669

8670-8679

8680-8689

8690-8699

8700-8709

8710-8719

8720-8729

8730-8739

8740-8749

8750-8759

8760-8769

8770-8779

8780-8789

8790-8799

8800-8809

8810-8819

8820-8829

8830-8839

8840-8849

8850-8859

8860-8869

8870-8879

8880-8889

8890-8899

8900-8909

8910-8919

8920-8929

8930-8939

8940-8949

8950-8959

8960-8969

8970-8979

8980-8989

8990-8999

9000-9009

9010-9019

9020-9029

9030-9039

9040-9049

9050-9059

9060-9069

9070-9079

9080-9089

9090-9099

9100-9109

9110-9119

9120-9129

9130-9139

9140-9149

9150-9159

9160-9169

9170-9179

9180-9189

9190-9199

9200-9209

9210-9219

9220-9229

9230-9239

9240-9249

9250-9259

9260-9269

9270-9279

9280-9289

9290-9299

9300-9309

9310-9319

9320-9329

9330-9339

9340-9349

9350-9359

9360-9369

9370-9379

9380-9389

9390-9399

9400-9409

9410-9419

9420-9429

9430-9439

9440-9449

9450-9459

9460-9469

9470-9479

9480-9489

9490-9499

9500-9509

9510-9519

9520-9529

9530-9539

9540-9549

9550-9559

9560-9569

9570-9579

9580-9589

9590-9599

9600-9609

9610-9619

9620-9629

9630-9639

9640-9649

9650-9659

9660-9669

9670-9679

9680-9689

9690-9699

9700-9709

9710-9719

9720-9729

9730-9739

9740-9749

9750-9759

9760-9769

9770-9779

9780-9789

9790-9799

9800-9809

9810-9819

9820-9829

9830-9839

9840-9849

9850-9859

9860-9869

9870-9879

9880-9889

9890-9899

9900-9909

9910-9919

992

10

Photochemical reaction of mercury vapors with oxygen. *Zhu, Fiklin.*

J. M. Frush, *J. Phys. Chem.* (U. S. S. R.) 8, 1013-30; *Acta Physicochim. U. R. S. S. I.* 8:30-64 (1964) (in German).

The oxidation of Hg vapors under the action of the resonance line 2537 Å. was observed at low pressures of O_2 (tenths of a mm.), as well as at pressures of several cm. The speed of reaction is proportional to the intensity of the line 2537 Å. and the no. of damped collisions with O_2 . At pressures of O_2 less than 1 mm. and with a 12-cm. absorbing column, the presence of light with wave lengths absorbed by O_2 (λ less than 2000 Å.) in the exciting beam has no influence on the speed of reaction. Whenever reaction takes place the formation of ozone is observed, but only in those parts of the vessel where the speed. HgO falls. At high pressures of O_2 , reaction occurs with wave lengths less than 2000 Å. Thus the presence of ozone is sufficient to cause reaction. The process of damping of the resonance line 2537 Å. by O_2 can be explained by the mechanism $Hg^* + O_2 \rightarrow Hg + O_2^*$, $O_2^* + O_2 \rightarrow O_3 + O$, $Hg + O_3 \rightarrow HgO + O_2$. At O formed in the second reaction either reacts directly with Hg in a triple collision or preliminarily unites with O_2 and forms ozone. The stability of the HgO mol. in the gas phase was not shown. It is possible that the reaction occurs only on the walls of the vessel. The 3 reactions fully explain the results of Dickinson and Sherrill obtained at high pressures (cf. *C. A.* 20, 1954). The mechanism $Hg^* + O_2 \rightarrow HgO + O$, $Hg^* + O_2 \rightarrow Hg' + O$, where Hg' is a metastable atom of Hg, is also possible. The part played by these processes in the reaction is not yet explained. Eino Hanninen

ASB-514 DETAILURGICAL LITERATURE CLASSIFICATION

TITLE AND SUB-TITLE		PROCESS AND PROPERTY INDEX		SUBJECT INDEX	
<div style="position: absolute; top: 10px; left: 10px; font-size: 2em;">S.2</div> <div style="position: absolute; top: 10px; right: 10px; font-size: 2em;">A 52</div> <div style="position: absolute; top: 300px; right: 10px; font-size: 1.2em;">D.C. AN 555</div> <div style="position: absolute; top: 300px; left: 10px; transform: rotate(-90deg); font-size: 0.8em;"> 2039. Observations of Night Sky Luminescence by an Extinction Method. N. Dobrotin, I. Frank and P. Cherenkov. <i>Comptes Rendus de l'Acad. des Sciences, U.S.S.R.</i> 1, pp. 110-117, Jan. 21, 1935. In English. Observations of the auroral green line 5577.3 Å in the light of the night sky have been made by the extinction method of Hennberg and Wawilow (see Abstract 4940 (1934)) using Rayleigh's auroral filter. It is found that the ratio of the intensity of the green line to the total intensity of the sky light varies from 4 % at the beginning of the night to 12 % at midnight. The absolute brightness of sky luminescence near the Pole star is found to be 1.5×10^{-6} candle per cm², giving for the intensity of the green line at the zenith at maximum brightness a value of 3.3×10^{-6} candle per cm². Evidence is brought forward against McLennan's view that the variation of intensity of the green line is due to quenching by visible light. [See following Abstract.] C. B. A. </div>		2039. Observations of Night Sky Luminescence by an Extinction Method. N. Dobrotin, I. Frank and P. Cherenkov. <i>Comptes Rendus de l'Acad. des Sciences, U.S.S.R.</i> 1, pp. 110-117, Jan. 21, 1935. In English. Observations of the auroral green line 5577.3 Å in the light of the night sky have been made by the extinction method of Hennberg and Wawilow (see Abstract 4940 (1934)) using Rayleigh's auroral filter. It is found that the ratio of the intensity of the green line to the total intensity of the sky light varies from 4 % at the beginning of the night to 12 % at midnight. The absolute brightness of sky luminescence near the Pole star is found to be 1.5×10^{-6} candle per cm ² , giving for the intensity of the green line at the zenith at maximum brightness a value of 3.3×10^{-6} candle per cm ² . Evidence is brought forward against McLennan's view that the variation of intensity of the green line is due to quenching by visible light. [See following Abstract.] C. B. A.		2039. Observations of Night Sky Luminescence by an Extinction Method. N. Dobrotin, I. Frank and P. Cherenkov. <i>Comptes Rendus de l'Acad. des Sciences, U.S.S.R.</i> 1, pp. 110-117, Jan. 21, 1935. In English. Observations of the auroral green line 5577.3 Å in the light of the night sky have been made by the extinction method of Hennberg and Wawilow (see Abstract 4940 (1934)) using Rayleigh's auroral filter. It is found that the ratio of the intensity of the green line to the total intensity of the sky light varies from 4 % at the beginning of the night to 12 % at midnight. The absolute brightness of sky luminescence near the Pole star is found to be 1.5×10^{-6} candle per cm ² , giving for the intensity of the green line at the zenith at maximum brightness a value of 3.3×10^{-6} candle per cm ² . Evidence is brought forward against McLennan's view that the variation of intensity of the green line is due to quenching by visible light. [See following Abstract.] C. B. A.	
ASD-51A METALLURGICAL LITERATURE CLASSIFICATION					
100000 01 100000 02 100000 03 100000 04 100000 05 100000 06 100000 07 100000 08 100000 09 100000 10 100000 11 100000 12 100000 13 100000 14 100000 15 100000 16 100000 17 100000 18 100000 19 100000 20 100000 21 100000 22 100000 23 100000 24 100000 25 100000 26 100000 27 100000 28 100000 29 100000 30 100000 31 100000 32 100000 33 100000 34 100000 35 100000 36 100000 37 100000 38 100000 39 100000 40 100000 41 100000 42 100000 43 100000 44 100000 45 100000 46 100000 47 100000 48 100000 49 100000 50 100000 51 100000 52 100000 53 100000 54 100000 55 100000 56 100000 57 100000 58 100000 59 100000 60 100000 61 100000 62 100000 63 100000 64 100000 65 100000 66 100000 67 100000 68 100000 69 100000 70 100000 71 100000 72 100000 73 100000 74 100000 75 100000 76 100000 77 100000 78 100000 79 100000 80 100000 81 100000 82 100000 83 100000 84 100000 85 100000 86 100000 87 100000 88 100000 89 100000 90 100000 91 100000 92 100000 93 100000 94 100000 95 100000 96 100000 97 100000 98 100000 99 100000 100		100000 01 100000 02 100000 03 100000 04 100000 05 100000 06 100000 07 100000 08 100000 09 100000 10 100000 11 100000 12 100000 13 100000 14 100000 15 100000 16 100000 17 100000 18 100000 19 100000 20 100000 21 100000 22 100000 23 100000 24 100000 25 100000 26 100000 27 100000 28 100000 29 100000 30 100000 31 100000 32 100000 33 100000 34 100000 35 100000 36 100000 37 100000 38 100000 39 100000 40 100000 41 100000 42 100000 43 100000 44 100000 45 100000 46 100000 47 100000 48 100000 49 100000 50 100000 51 100000 52 100000 53 100000 54 100000 55 100000 56 100000 57 100000 58 100000 59 100000 60 100000 61 100000 62 100000 63 100000 64 100000 65 100000 66 100000 67 100000 68 100000 69 100000 70 100000 71 100000 72 100000 73 100000 74 100000 75 100000 76 100000 77 100000 78 100000 79 100000 80 100000 81 100000 82 100000 83 100000 84 100000 85 100000 86 100000 87 100000 88 100000 89 100000 90 100000 91 100000 92 100000 93 100000 94 100000 95 100000 96 100000 97 100000 98 100000 99 100000 100			

ca

3

Coherent visible radiation of fast electrons passing through matter. I. Frank and Ig. Tamm. *Compt. rend. acad. sci. U. R. S. S. 16*, 109-14 (1937) (In English).

The phenomenon of the peculiar visible radiation from liquids and solids bombarded by fast electrons, such as β -electrons or Compton electrons, can be explained if the fact is considered that an electron moving in a medium radiates light even if it is moving uniformly provided its velocity is greater than the velocity of light in the medium. Seventeen equations are derived in developing this theory.

M. McMahon

Doc. AN JSSR

ASAC-3.4 METALLURGICAL LITERATURE CLASSIFICATION

Visible radiation of pure liquids under the action of rapid electrons. I. E. Tamm, L. M. Erank and P. A. Cherenkov. *Bull. acad. sci. U. R. S. S. Div. sci. math nat., Ser. phys.* 1938, 29, 30 (in English 30 1). On passing rapid Compton electrons from γ -rays or β -particles from radioactive substances through liquids, a hitherto unknown faint luminescence occurs which is difficult to observe because of the strong ordinary fluorescence. A careful investigation revealed the following fundamental properties of this luminescence. Unlike radiation produced by γ -rays, the intensity of this new radiation produced by rapid electrons is inversely proportional to the density of the liquid. It is not quenched by change in viscosity produced by heating of the liquid or by dissolving in it such salts as KI or AgNO_3 . It is partly polarized and the degree of polarization is affected by change of viscosity of the liquid; it is in the direction of motion of the electrons and produces a continuous spectrum. S. I. Madorsky

Vest. 111 1004, 1004

ASO-51A METALLURGICAL LITERATURE CLASSIFICATION

Pair formation in nitrogen by γ rays I. V. Gushchik and A. M. Frank. *Bull. Acad. Sci. U. R. S. S. (Chem. sec. math. nat., Ser. phys. 1938, No. 5, 673) in English, 703-4; cf. C. A. 33, 2429.*—The formation of electron-positron pairs by Th^{γ} - γ rays was studied in an air-filled Wilson chamber. There were 32 pairs on 1930 stereoscopic photographs. The energies of 28 pairs were determined from the curvatures of their paths in a magnetic field. A sharp max. was found for pairs with 1.0 m. e. v. Out of 24 pairs with energies of more than 1 m. e. v., 10 pairs had energies from 1.45 to 1.75 m. e. v., and 12 pairs had energies from 1.65 to 1.65 m. e. v. The av. energy of the positrons for the obtained pairs was considerably higher than that of the electrons. Their difference $E_+ - E_-$ was from 0.2 to 0.3 m. e. v. while the calculated difference was 0.02 m. e. v. The cross section for pair formation was $0.9 \times 10^{-28} \text{ cm}^2$ instead of $1.8 \times 10^{-28} \text{ cm}^2$, which is expected theoretically. One of the reasons for this deviation was that the pairs whose energies were not measured were not taken into consideration. The comparison of the cross sections for pair-formation in N with the values for Kr obtained previously ($0.23 \times 10^{-28} \text{ cm}^2$) gives a value of their ratio which corresponds to the ratio $Z_1^2/Z_2^2 = 27$, where Z is the at. no. Munich

13. 12. 1899

1ST AND 2ND ORDERS										100 AND 4TH ORDERS									
PROCESSES AND PROPERTIES INDEX																			
<p>SA</p> <p>2647. Probability of Pair Creation in Kr by γ-Rays. L. V. Groshove and I. M. Frank. <i>Comptes Rendus (Doklady) de l'Acad. des Sciences, U.S.S.R.</i> 18. 7. pp. 419-422, 1938. In English.—Using a special hermetically sealed cloud chamber, the creation of pairs of positrons and electrons in Kr by the γ-rays from Rd-Th is examined. The results are divided into two groups covering the energy intervals 0.8-1 eMV and greater than 1 eMV respectively. Methods are devised for allowing for spurious pairs due to electrons which have suffered single nuclear scattering. The effective cross-section for pair production is found to be in approximate agreement with the theory of Jaeger and Hulme. [See following Abstract.] F. C. C.</p> <p>45366</p> <p>DAL-AN 551K</p>																			
ASB-5LA METALLURGICAL LITERATURE CLASSIFICATION																			
10000 100000										10000 100000									
10000 100000										10000 100000									

1ST AND 2ND CODES										3RD AND 4TH CODES									
PROCESSES AND PROPERTIES INDEX																			
1A										A5366									
<p>2648. Angular Distribution of Pairs in Kr. L. V. Grodzov and I. M. Frank. <i>Comptes Rendus (Doklady) de l'Acad. des Sciences, U.S.S.R.</i> 19. 1-8. pp. 40-51, 1966. In English.—Continuing previous work [see preceding Abstract] examination is made of the angular distribution of pairs in Kr produced by γ-rays from ThC''. The data agree with the theory of Bethe and Heitler. There appears to be a real difference in the angular distribution of the positrons and electrons, particularly at angles $< 10^\circ$, for this region includes 84% of the positrons and only 18% of the electrons. The mean energy difference between the positrons and electrons is 100 eV and is in agreement with theory. This difference increases with increasing atomic number. The average angle between the pairs is 46°, which is considerably higher than that reported by previous workers. [See following Abstract.] F. C. C.</p>																			
<p>ASAC-36.0 METALLURGICAL LITERATURE CLASSIFICATION</p>																			
1ST AND 2ND CODES										3RD AND 4TH CODES									
1ST AND 2ND CODES										3RD AND 4TH CODES									

3

Nuclear impulse in pair creation. I. V. Goushev and I. M. Frank. *Compt. rend. acad. Sci. U. R. S. S.* 19, 280-42 (1968) (in English). Previous articles of I. V. Goushev and I. M. Frank, cf. C. A. 32, 6146, 6041, considered the angular distribution and probability of pair production in Kr by γ -rays of Th C². The impulse transmitted to the nucleus during this process is now studied. In the act of pair creation practically no energy is transmitted to the nucleus but a certain impulse P is always transmitted. The av. value of P calculated for 20 pairs is 1.7 μ c. The angle ϕ between photon impulse and that of the pair was not large, for the 20 pairs the av. value of ϕ is 17°. Three pairs in which the positron was carrying the full energy were omitted in analyzing the impulses.

M. McMahon

ASD-55A METALLURGICAL LITERATURE CLASSIFICATION

PROCESSES AND PROPERTIES INDEX

Production of pairs in nitrogen by γ -rays. I. V. Groshev and I. M. Frank. *Compt. rend. acad. sci. U. R. S. S.* 20, 373-8 (1960) (in English); cf. C. A. B., 4649. - A sharp max. in the no. of pairs appears at 1.0 m. e. v. In most cases the positrons have a greater energy than the electrons. Benn's approximation cor. for Z would account for this.
G. M. Evans

ASTM-SLA METALLURGICAL LITERATURE CLASSIFICATION

A new type of nuclear reactions (The splitting of uranium and thorium nuclei under the influence of neutrons). I. M. Frank, *Pravda*, No. 22, 20, 1940, 1941, 1942, 1943, 1944, 1945, 1946, 1947, 1948, 1949, 1950, 1951, 1952, 1953, 1954, 1955, 1956, 1957, 1958, 1959, 1960, 1961, 1962, 1963, 1964, 1965, 1966, 1967, 1968, 1969, 1970, 1971, 1972, 1973, 1974, 1975, 1976, 1977, 1978, 1979, 1980, 1981, 1982, 1983, 1984, 1985, 1986, 1987, 1988, 1989, 1990, 1991, 1992, 1993, 1994, 1995, 1996, 1997, 1998, 1999, 2000, 2001, 2002, 2003, 2004, 2005, 2006, 2007, 2008, 2009, 2010, 2011, 2012, 2013, 2014, 2015, 2016, 2017, 2018, 2019, 2020, 2021, 2022, 2023, 2024, 2025, 2026, 2027, 2028, 2029, 2030, 2031, 2032, 2033, 2034, 2035, 2036, 2037, 2038, 2039, 2040, 2041, 2042, 2043, 2044, 2045, 2046, 2047, 2048, 2049, 2050, 2051, 2052, 2053, 2054, 2055, 2056, 2057, 2058, 2059, 2060, 2061, 2062, 2063, 2064, 2065, 2066, 2067, 2068, 2069, 2070, 2071, 2072, 2073, 2074, 2075, 2076, 2077, 2078, 2079, 2080, 2081, 2082, 2083, 2084, 2085, 2086, 2087, 2088, 2089, 2090, 2091, 2092, 2093, 2094, 2095, 2096, 2097, 2098, 2099, 2100, 2101, 2102, 2103, 2104, 2105, 2106, 2107, 2108, 2109, 2110, 2111, 2112, 2113, 2114, 2115, 2116, 2117, 2118, 2119, 2120, 2121, 2122, 2123, 2124, 2125, 2126, 2127, 2128, 2129, 2130, 2131, 2132, 2133, 2134, 2135, 2136, 2137, 2138, 2139, 2140, 2141, 2142, 2143, 2144, 2145, 2146, 2147, 2148, 2149, 2150, 2151, 2152, 2153, 2154, 2155, 2156, 2157, 2158, 2159, 2160, 2161, 2162, 2163, 2164, 2165, 2166, 2167, 2168, 2169, 2170, 2171, 2172, 2173, 2174, 2175, 2176, 2177, 2178, 2179, 2180, 2181, 2182, 2183, 2184, 2185, 2186, 2187, 2188, 2189, 2190, 2191, 2192, 2193, 2194, 2195, 2196, 2197, 2198, 2199, 2200, 2201, 2202, 2203, 2204, 2205, 2206, 2207, 2208, 2209, 2210, 2211, 2212, 2213, 2214, 2215, 2216, 2217, 2218, 2219, 2220, 2221, 2222, 2223, 2224, 2225, 2226, 2227, 2228, 2229, 2230, 2231, 2232, 2233, 2234, 2235, 2236, 2237, 2238, 2239, 2240, 2241, 2242, 2243, 2244, 2245, 2246, 2247, 2248, 2249, 2250, 2251, 2252, 2253, 2254, 2255, 2256, 2257, 2258, 2259, 2260, 2261, 2262, 2263, 2264, 2265, 2266, 2267, 2268, 2269, 2270, 2271, 2272, 2273, 2274, 2275, 2276, 2277, 2278, 2279, 2280, 2281, 2282, 2283, 2284, 2285, 2286, 2287, 2288, 2289, 2290, 2291, 2292, 2293, 2294, 2295, 2296, 2297, 2298, 2299, 2300, 2301, 2302, 2303, 2304, 2305, 2306, 2307, 2308, 2309, 2310, 2311, 2312, 2313, 2314, 2315, 2316, 2317, 2318, 2319, 2320, 2321, 2322, 2323, 2324, 2325, 2326, 2327, 2328, 2329, 2330, 2331, 2332, 2333, 2334, 2335, 2336, 2337, 2338, 2339, 2340, 2341, 2342, 2343, 2344, 2345, 2346, 2347, 2348, 2349, 2350, 2351, 2352, 2353, 2354, 2355, 2356, 2357, 2358, 2359, 2360, 2361, 2362, 2363, 2364, 2365, 2366, 2367, 2368, 2369, 2370, 2371, 2372, 2373, 2374, 2375, 2376, 2377, 2378, 2379, 2380, 2381, 2382, 2383, 2384, 2385, 2386, 2387, 2388, 2389, 2390, 2391, 2392, 2393, 2394, 2395, 2396, 2397, 2398, 2399, 2400, 2401, 2402, 2403, 2404, 2405, 2406, 2407, 2408, 2409, 2410, 2411, 2412, 2413, 2414, 2415, 2416, 2417, 2418, 2419, 2420, 2421, 2422, 2423, 2424, 2425, 2426, 2427, 2428, 2429, 2430, 2431, 2432, 2433, 2434, 2435, 2436, 2437, 2438, 2439, 2440, 2441, 2442, 2443, 2444, 2445, 2446, 2447, 2448, 2449, 2450, 2451, 2452, 2453, 2454, 2455, 2456, 2457, 2458, 2459, 2460, 2461, 2462, 2463, 2464, 2465, 2466, 2467, 2468, 2469, 2470, 2471, 2472, 2473, 2474, 2475, 2476, 2477, 2478, 2479, 2480, 2481, 2482, 2483, 2484, 2485, 2486, 2487, 2488, 2489, 2490, 2491, 2492, 2493, 2494, 2495, 2496, 2497, 2498, 2499, 2500, 2501, 2502, 2503, 2504, 2505, 2506, 2507, 2508, 2509, 2510, 2511, 2512, 2513, 2514, 2515, 2516, 2517, 2518, 2519, 2520, 2521, 2522, 2523, 2524, 2525, 2526, 2527, 2528, 2529, 2530, 2531, 2532, 2533, 2534, 2535, 2536, 2537, 2538, 2539, 2540, 2541, 2542, 2543, 2544, 2545, 2546, 2547, 2548, 2549, 2550, 2551, 2552, 2553, 2554, 2555, 2556, 2557, 2558, 2559, 2560, 2561, 2562, 2563, 2564, 2565, 2566, 2567, 2568, 2569, 2570, 2571, 2572, 2573, 2574, 2575, 2576, 2577, 2578, 2579, 2580, 2581, 2582, 2583, 2584, 2585, 2586, 2587, 2588, 2589, 2590, 2591, 2592, 2593, 2594, 2595, 2596, 2597, 2598, 2599, 2600, 2601, 2602, 2603, 2604, 2605, 2606, 2607, 2608, 2609, 2610, 2611, 2612, 2613

W. R. Hens

ALPHABETIC INDEX																									
A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	Q	R	S	T	U	V	W	X	Y	Z
<p>PHYSICAL PROPERTIES INDEX</p> <p>Doppler effect in a refractive medium. I. M. Frank. <i>Zh. Fiz.</i> 3 <i>J. Phys. (U. S. S. R.)</i> 7, 40-67 (1943) (in English). Theoret.—mathematical. A "complex Doppler effect" can be observed experimentally for velocities much less than that of light in a gas possessing a sufficiently narrow absorption line. Cherenkov type radiation is emitted if the dipole moment of the source has a const. component. P. H. Rathmann</p>																									
<p>ASTRO-SEA METALLURGICAL LITERATURE CLASSIFICATION</p> <p>ASTRO-SEA METALLURGICAL LITERATURE CLASSIFICATION</p>																									

FRANK, I. M.

"Radiation Receiver with Reduced Sensitivity to the Position of the Source,"
Dokl. AN USSR, 39, No 2, 1943. Physics Inst. im. Lebedev, AS USSR

FRANK, I. M.

"Interference Phenomena in the Case of Cherenkov Radiation," Dok AN USSR, 42, No.8,
1943. Phys. Inst. im. Lebedev, AS USSR

1ST AND 2ND ORDER

PROCESSES AND PROPERTIES INDEX

3

CA

Radiation of electron upon uniform motion in a refracting medium. I. E. Tamm and I. M. Frank. *Trudy Fiz. Inst. im. P. N. Lebedeva, Akad. Nauk S. S. S. R.* 2, No. 4, 63-B(1944).—Math.-theoret. In the Cherenkov radiation produced by fast electrons passing through matter, the effect is a consequence of Sommerfeld's considerations (*Götting. Nachricht.* 90, 363(1904)) of super-light-speed electrons. The qual. and quant. properties of the radiation are explainable on the basis of the electron velocity exceeding the phase velocity of light, which is a real condition in suvraming media, e. g., H_2O . G. M. K.

ASB-SEA METALLURGICAL LITERATURE CLASSIFICATION

SECTION 1

SECTION 2

SECTION 3

SECTION 4

SECTION 5

SECTION 6

SECTION 7

SECTION 8

SECTION 9

SECTION 10

SECTION 11

SECTION 12

SECTION 13

SECTION 14

SECTION 15

SECTION 16

SECTION 17

SECTION 18

SECTION 19

SECTION 20

SECTION 21

SECTION 22

SECTION 23

SECTION 24

SECTION 25

SECTION 26

SECTION 27

SECTION 28

SECTION 29

SECTION 30

SECTION 31

SECTION 32

SECTION 33

SECTION 34

SECTION 35

SECTION 36

SECTION 37

SECTION 38

SECTION 39

SECTION 40

SECTION 41

SECTION 42

SECTION 43

SECTION 44

SECTION 45

SECTION 46

SECTION 47

SECTION 48

SECTION 49

SECTION 50

SECTION 51

SECTION 52

SECTION 53

SECTION 54

SECTION 55

SECTION 56

SECTION 57

SECTION 58

SECTION 59

SECTION 60

SECTION 61

SECTION 62

SECTION 63

SECTION 64

SECTION 65

SECTION 66

SECTION 67

SECTION 68

SECTION 69

SECTION 70

SECTION 71

SECTION 72

SECTION 73

SECTION 74

SECTION 75

SECTION 76

SECTION 77

SECTION 78

SECTION 79

SECTION 80

SECTION 81

SECTION 82

SECTION 83

SECTION 84

SECTION 85

SECTION 86

SECTION 87

SECTION 88

SECTION 89

SECTION 90

SECTION 91

SECTION 92

SECTION 93

SECTION 94

SECTION 95

SECTION 96

SECTION 97

SECTION 98

SECTION 99

SECTION 100

BC

Radiation of a uniformly moving electron due to its transition from one medium into another. V. G. Glazov and V. Glaznov (J. Phys. U.S.S.R., 1948, 8, No. 10, 1645, 1646, 1647, 1648, 1649, 1650, 1651, 1652, 1653, 1654, 1655, 1656, 1657, 1658, 1659, 1660, 1661, 1662, 1663, 1664, 1665, 1666, 1667, 1668, 1669, 1670, 1671, 1672, 1673, 1674, 1675, 1676, 1677, 1678, 1679, 1680, 1681, 1682, 1683, 1684, 1685, 1686, 1687, 1688, 1689, 1690, 1691, 1692, 1693, 1694, 1695, 1696, 1697, 1698, 1699, 1700, 1701, 1702, 1703, 1704, 1705, 1706, 1707, 1708, 1709, 1710, 1711, 1712, 1713, 1714, 1715, 1716, 1717, 1718, 1719, 1720, 1721, 1722, 1723, 1724, 1725, 1726, 1727, 1728, 1729, 1730, 1731, 1732, 1733, 1734, 1735, 1736, 1737, 1738, 1739, 1740, 1741, 1742, 1743, 1744, 1745, 1746, 1747, 1748, 1749, 1750, 1751, 1752, 1753, 1754, 1755, 1756, 1757, 1758, 1759, 1760, 1761, 1762, 1763, 1764, 1765, 1766, 1767, 1768, 1769, 1770, 1771, 1772, 1773, 1774, 1775, 1776, 1777, 1778, 1779, 1780, 1781, 1782, 1783, 1784, 1785, 1786, 1787, 1788, 1789, 1790, 1791, 1792, 1793, 1794, 1795, 1796, 1797, 1798, 1799, 1800, 1801, 1802, 1803, 1804, 1805, 1806, 1807, 1808, 1809, 1810, 1811, 1812, 1813, 1814, 1815, 1816, 1817, 1818, 1819, 1820, 1821, 1822, 1823, 1824, 1825, 1826, 1827, 1828, 1829, 1830, 1831, 1832, 1833, 1834, 1835, 1836, 1837, 1838, 1839, 1840, 1841, 1842, 1843, 1844, 1845, 1846, 1847, 1848, 1849, 1850, 1851, 1852, 1853, 1854, 1855, 1856, 1857, 1858, 1859, 1860, 1861, 1862, 1863, 1864, 1865, 1866, 1867, 1868, 1869, 1870, 1871, 1872, 1873, 1874, 1875, 1876, 1877, 1878, 1879, 1880, 1881, 1882, 1883, 1884, 1885, 1886, 1887, 1888, 1889, 1890, 1891, 1892, 1893, 1894, 1895, 1896, 1897, 1898, 1899, 1900, 1901, 1902, 1903, 1904, 1905, 1906, 1907, 1908, 1909, 1910, 1911, 1912, 1913, 1914, 1915, 1916, 1917, 1918, 1919, 1920, 1921, 1922, 1923, 1924, 1925, 1926, 1927, 1928, 1929, 1930, 1931, 1932, 1933, 1934, 1935, 1936, 1937, 1938, 1939, 1940, 1941, 1942, 1943, 1944, 1945, 1946, 1947, 1948, 1949, 1950, 1951, 1952, 1953, 1954, 1955, 1956, 1957, 1958, 1959, 1960, 1961, 1962, 1963, 1964, 1965, 1966, 1967, 1968, 1969, 1970, 1971, 1972, 1973, 1974, 1975, 1976, 1977, 1978, 1979, 1980, 1981, 1982, 1983, 1984, 1985, 1986, 1987, 1988, 1989, 1990, 1991, 1992, 1993, 1994, 1995, 1996, 1997, 1998, 1999, 2000, 2001, 2002, 2003, 2004, 2005, 2006, 2007, 2008, 2009, 2010, 2011, 2012, 2013, 2014, 2015, 2016, 2017, 2018, 2019, 2020, 2021, 2022, 2023, 2024, 2025, 2026, 2027, 2028, 2029, 2030, 2031, 2032, 2033, 2034, 2035, 2036, 2037, 2038, 2039, 2040, 2041, 2042, 2043, 2044, 2045, 2046, 2047, 2048, 2049, 2050, 2051, 2052, 2053, 2054, 2055, 2056, 2057, 2058, 2059, 2060, 2061, 2062, 2063, 2064, 2065, 2066, 2067, 2068, 2069, 2070, 2071, 2072, 2073, 2074, 2075, 2076, 2077, 2078, 2079, 2080, 2081, 2082, 2083, 2084, 2085, 2086, 2087, 2088, 2089, 2090, 2091, 2092, 2093, 2094, 2095, 2096, 2097, 2098, 2099, 2100, 2101, 2102, 2103, 2104, 2105, 2106, 2107, 2108, 2109, 2110, 2111, 2112, 2113, 2114, 2115, 2116, 2117, 2118, 2119, 2120, 2121, 2122, 2123, 2124, 2125, 2126, 2127, 2128, 2129, 2130, 2131, 2132, 2133, 2134, 2135, 2136, 2137, 2138, 2139, 2140, 2141, 2142, 2143, 2144, 2145, 2146, 2147, 2148, 2149, 2150, 2151, 2152, 2153, 2154, 2155, 2156, 2157, 2158, 2159, 2160, 2161, 2162, 2163, 2164, 2165, 2166, 2167, 2168, 2169, 2170, 2171, 2172, 2173, 2174, 2175, 2176, 2177, 2178, 2179, 2180, 2181, 2182, 2183, 2184, 2185, 2186, 2187, 2188, 2189, 2190, 2191, 2192, 2193, 2194, 2195, 2196, 2197, 2198, 2199, 2200, 2201, 2202, 2203, 2204, 2205, 2206, 2207, 2208, 2209, 2210, 2211, 2212, 2213, 2214, 2215, 2216, 2217, 2218, 2219, 2220, 2221, 2222, 2223, 2224, 2225, 2226, 2227, 2228, 2229, 2230, 2231, 2232, 2233, 2234, 2235, 2236, 2237, 2238, 2239, 2240, 2241, 2242, 2243, 2244, 2245, 2246, 2247, 2248, 2249, 2250, 2251, 2252, 2253, 2254, 2255, 2256, 2257, 2258, 2259, 2260, 2261, 2262, 2263, 2264, 2265, 2266, 2267, 2268, 2269, 2270, 2271, 2272, 2273, 2274, 2275, 2276, 2277, 2278, 2279, 2280, 2281, 2282, 2283, 2284, 2285, 2286, 2287, 2288, 2289, 2290, 2291, 2292, 2293, 2294, 2295, 2296, 2297, 2298, 2299, 2300, 2301, 2302, 2303, 2304, 2305, 2306, 2307, 2308, 2309, 2310, 2311, 2312, 2313, 2314, 2315, 2316, 23

Zhou, F. et al.

H. R. C.

A 30 3 L A METALLURGICAL LITERATURE CLASSIFICATION

CLASSIFICATION		PROCESS AND PROPERTIES INDEX		SUBJECTS	
C-1					
<p>Compensation chamber for analyzing the composition of radioactive preparations with respect to γ-rays. I. Frank. <i>J. Exptl. Theoret. Phys.</i> (U.S.S.R.) 18, 25-31(1948) (English summary); <i>J. Phys.</i> (U.S.S.R.) 9, 123-9.—The ionization current due to the γ-rays was measured by means of 3 ionization chambers arranged in a compensation circuit. The walls of the chambers consisted of successive layers a, b, c in the first and a, c, b in the second case, where a and b are the substances with small atomic numbers (as e.g., Al), and c is Pb. This method of compensation chambers permits: (1) measurement of the wave lengths of monochromatic γ-rays provided the apparatus is previously calibrated, (2) demonstration of the complexity of γ-rays spectrum, and (3) detn. of the content of Ra, Rd-Th, and Ms-Th in radioactive samples. R. C.</p>					
Leningrad Phys. Inst., AS USSR					
ASB-SLA METALLURGICAL LITERATURE CLASSIFICATION					
SOURCE SYNONYMS		SYNOPSIS WITH ONLY ONE		ORIGINATOR	
SOURCE NO.		SYNOPSIS NO.		ORIGINATOR NO.	
1		1		1	
2		2		2	
3		3		3	
4		4		4	
5		5		5	
6		6		6	
7		7		7	
8		8		8	
9		9		9	
10		10		10	
11		11		11	
12		12		12	
13		13		13	
14		14		14	
15		15		15	
16		16		16	
17		17		17	
18		18		18	
19		19		19	
20		20		20	
21		21		21	
22		22		22	
23		23		23	
24		24		24	
25		25		25	
26		26		26	
27		27		27	
28		28		28	
29		29		29	
30		30		30	
31		31		31	
32		32		32	
33		33		33	
34		34		34	
35		35		35	
36		36		36	
37		37		37	
38		38		38	
39		39		39	
40		40		40	
41		41		41	
42		42		42	
43		43		43	
44		44		44	
45		45		45	
46		46		46	
47		47		47	
48		48		48	
49		49		49	
50		50		50	
51		51		51	
52		52		52	
53		53		53	
54		54		54	
55		55		55	
56		56		56	
57		57		57	
58		58		58	
59		59		59	
60		60		60	
61		61		61	
62		62		62	
63		63		63	
64		64		64	
65		65		65	
66		66		66	
67		67		67	
68		68		68	
69		69		69	
70		70		70	
71		71		71	
72		72		72	
73		73		73	
74		74		74	
75		75		75	
76		76		76	
77		77		77	
78		78		78	
79		79		79	
80		80		80	
81		81		81	
82		82		82	
83		83		83	
84		84		84	
85		85		85	
86		86		86	
87		87		87	
88		88		88	
89		89		89	
90		90			

The transition effect of γ rays and their influence on the ionization current in ionization chambers. O. N. Vavilov and I. M. Frank. *Zhur. Exptl. Teoret. Fis.* 17, 176-87 (1947); *J. Phys. (U.S.S.R.)* 11, No. 3(1947) (in English).

The substances to be studied were placed above or below a $40 \times 40 \times 5$ cm. chamber with 0.015 mm Al walls in such a way that the total thickness of each substance penetrated by the incident radiation remained constant. The ionization current I was studied as a function of x_1 , the thickness of the lighter material next to the chamber. When the filters were between the γ -source and the chamber, the Fe-Al, Al-C, and Fe-C curves rose to a plateau at $x_1 \leq 1$ g. cm 2 ; the Pb-Al, Pb-Fe, and Pb-C curves showed sharp min. at $x_1 \approx 0.15$ g. cm 2 . The curves are accounted for in terms of Compton effect and photoeffect. Curves are also given for the cases in which the filters were placed on the other side of the chamber, and on both sides. The following values were obtained for I_0 , the ionization current due to radiation having a component in the sense of the incident radiation, and I_s , that due to oppositely directed radiation. The chamber current caused by electrons resulting from the ionization of air was taken as unity. For C, Al, Fe, and Pb, $I_0 = 2.30, 2.19, 2.02$, and 2.02 , resp., and $I_s = 0.16, 0.48, 0.78$, and 2.02 , resp.

Cyrus Feilman

Cyrus Feldman

USSR/Nuclear Phys - Gamma Rays

Nuclear Phys - Impact, Electronic

Feb 1947

"Angular Distribution of Electronic Pairs Produced by Gamma Rays of ThC," L. V. Groshev, I. M. Frank, Phys Inst imeni P. M. Lebedev, Acad Sci USSR, 2¹/₂ pp

"Zhur Eksper i Teoret Fiz" Vol XVII, No 2

Shows that dependence of distribution of angles between components of pairs formed by Gamma rays on the atomic number of the irradiated substance can differ, depending on method of pair registration employed. As result of this, the difference in results obtained with use of the Wilson chamber and the counters does not lead to contradiction.

FA 57T70

FRANK, I.M. and CINZBERG, V.L.


"The Doppler Effect at Super-phase Velocity." Dok. Akad. Nauk. SSSR,
56 (1947), 6, 583-586.

SO: Translation— 30 Apr 1954.

CA

The radiation from electrons and atoms which are in motion along the axis of a canal in a dense medium. V. L. Ginsburg and I. M. Frank. *Doklady Akad. Nauk S.S.S.R.* 99, 699-702 (1947); *Chem. Zentr.* (Russian Zone Ed.) 1948, 11, 138; cf. C.A. 43, 11156. — A theoretical and math. discussion is presented to show that it is possible for Cherenkov radiation to be produced by an electron moving within a canal of small radius ($r \ll \lambda$) and that it is further possible for the Doppler effect to be produced at velocities greater than that of light. The simplest case of the motion of an electron and of an atom along a canal at a velocity $v = \beta c$ is considered. If $n_1 = n_2$ (inside and outside the canal) the Cherenkov effect would be produced. It is shown that the condition $\beta < 5 \times 10^{-2}$ is not necessary for the development of this effect. The radiation of an oscillator moving with uniform velocity can be considered in exactly the same manner as that of an electron. The Doppler effect is theoretically possible at velocities greater than that of light if the electron moves along the boundary medium-vacuum at a distance $\approx 0.1 \lambda$ from the medium. M. G. Moore

Translation - 2524467. 30 Apr 57



Quenching of fluorescence by a light-absorbing medium.
M. D. Galanin and I. M. Frank (P.N. Lebedev Phys. Inst.
Acad. Sci. U.S.S.R., Moscow). *Zhur. Fiz. i Tverd. Fis.*
21, 114-90 (1951); cf. preceding abstr.—Consideration, by
classical electrodynamics, of a model consisting of an oscillat-
ing dipole of moment p , emitting a frequency ω , and placed
in a homogeneous medium absorbing that frequency, gives
in the case of $\omega\lambda/2\pi \ll 1$, for the energy flux at a distance
 $R \sim \lambda/2\pi$, the expression $S_0 = (1/4\pi) p^2 \omega^2 / c^3$, i.e. the same
as for a transparent medium; at greater R , S decreases ac-
cording to $S = S_0 \cdot \exp(-\alpha R)$, and within the sphere $R \ll \lambda/2\pi$, an
addnl. energy is absorbed in the amt. of $S_0 \omega (\lambda/2\pi)^3$, where α
is an empirical const., greater than the sum of the radii of the
interacting mole., and representing the min. R . Corre-
spondingly, in the absorbing medium, the mean life τ of the
excited state becomes $1/\tau = (1/\tau_0) [1 + (\alpha\lambda/2\pi)(\lambda/2\pi)^3]$,
i.e. the relative variation of τ_0 is independent of its value, in
contrast to ordinary collision quenching. In application to
real fluorescent solns., p is a function of λ . On the assump-
tion that the form of the function $p(\lambda)$ does not change as a
result of absorption, the energy given up in an absorbing
medium is $W = \int_0^\infty S(\lambda) [1 + (\alpha(\lambda)(\lambda/2\pi)^3/a^3)] d\lambda \approx W_0 +$
 ΔW , and the variation of τ should be of the form $1/\tau =$
 $1/\tau_0 [1 + (\Delta W/W_0)]$, where $\Delta W = \int_0^\infty (1/a^3) S(\lambda)(\lambda/2\pi)^3$
 $\alpha(\lambda) d\lambda$ and $W_0 = \int_0^\infty S(\lambda) d\lambda$. The quenching is thus detd. by
an overlap of the fluorescence and absorption spectra $S(\lambda)$ -
 $\alpha(\lambda)$ and the factor $(\lambda/2\pi)^3$, and this result is in approx.
accord with exptl. observations (cf. following abstr., Part
II) with $a \sim 1 \times 10^{-10}$ cm. Similar developments are given
for a magnetic dipole and an elec. quadrupole. N. T.

FRANK, I. M., GAFIRO, A. L., SHIRANIK, A. V., ANTONOV, A. I.,
KORIN, I. D., and NEUFODYEV, B. A.

"A Study of Neutron Diffusion in Beryllium Graphite and water by the
Impulse method," a paper presented at the Atoms for Peace Conference,
Geneva, Switzerland, 1955

FRANK, I. M., FEYNBERG, Ye. L., BROSHOV, L. V., SHAIRO, F. L., SHTRAVENH, I. V.,
KOZINETZ, O. I., LAZAREVA, L. Ye. and TOLSTOV, K. D.

"Investigation of the Parameters of Uranium-Graphite Systems by the Prism Method".

Report appearing in 1st Volume of "Session of the Academy of Sciences USSR On the Peaceful use of Atomic Energy, 1-5 July 1955", Publishing House of Sciences USSR, 1955.

SO: Sum 728, 28 Nov 1955.

FRANK, I.M.

USSR
4

The multiplication of neutrons in uranium-graphite systems. I. V. Groshev, E. I. Feinberg, and I. M. Frank. *Soviet Acad. Nauk S.S.S.R. po Mirnomu Tipu i Energii Atomnoi Energii, Zasedaniya Otdel. Fiz.-Mat. Nauk* 1955, 3-18 (English summary, 19-20); cf. following 3 abstracts. —The phys. aspect of neutron multiplication in a heterogeneous U-graphite system was discussed from the exptl. and theoretical standpoint. The expts. were carried out by the exponential method by using various consens. of U and for various temp. conditions. The values of the thermal neutron utilization const., θ , and of the multiplication const., k , were detd., and the effect of the air gap and of the water jacket around the slugs was studied. J. Rovtar Leach.

1 RAM

Physic Inst. in P.N. Lebedev, AS USSR 7m

FRANK, I. D.

Measurement of temperature effects in uranium-graphite subcritical systems. B. P. Ad'yasevich, O. I. Koznets, K. D. Toksov, I. M. Frank, P. L. Shapiro, and I. V. Shtrankh. *Sessiya Akad. Nauk S.S.S.R. po Mirnomu Ispol'zovaniyu Atomnoi Energii, Zasedaniya Otdel. Fiz.-Mat. Nauk* 1955, 132-54 (English summary, 154-5).—The temp. effects on the multiplication const. $k_{\infty} = \eta \epsilon \theta$ and also on all the factors thereof were investigated by two $120 \times 120 \times 260$ cm. prisms, each in its own thermostat, heated to $300-450^{\circ}$. U-graphite (I) lattices were used with slug diams. of 32-7 mm. with various U concns. (c). A Ra-Be and a Po-Be neutron source were employed. The thermal utilization factor θ was measured by the C1 ratio method; a pos. temp. effect was found which increased with decreasing U concn. and was greater in the presence of cooling H_2O . This effect was already predicted by the elementary diffusion theory, but the theory furnished too high values in the absence and too low values in the presence of H_2O . This discrepancy was due to the fact that θ increased when the neutrons were cooled in H_2O upon entering the slug. The pos. effect on θ increased also if the cooling was extended over a layer of I adjacent to the slug. At identical U concns. the θ of a heterogeneous system can be increased over that of a homogeneous one by inhomogeneously heating the moderator. In order to evaluate correctly the temp. effect on the ϵ escape probability, ϵ , cold H_2O and H_2O at 80° were circulated alternately through the slugs, the temp. of the I being kept const. The variation of ϵ was obtained from exponential measurements of the buckling χ^2 and from measurements of the influence of heating on the epi-C1 neutron d. in the vicinity of the source. The temp. effect of the ϵ -integral is found to be $\partial \epsilon / \partial t = (1.95 \pm 0.4) \times 10^{-4}$ per degree. The temp. effect on k_{∞} was detd. by measuring χ^2 by the exponential method. The temp. changes of η (no. of neutrons generated/no. of neutrons captured) were calcd. from the changes of ϵ and θ and from the temp. effect on k_{∞} . It was found that η has a neg. temp. effect, which is roughly proportional to the variation of the mean energy of the thermal neutrons, caused by the heating of the system: $\partial \eta / \partial E = - (37 \pm 8) \% / e.v.$ Werner Jacobson

FRANK, I.M.
USSR/Nuclear Physics - Fission of U by negative pi-mesons

FD-2349

Card 1/2

Pub. 146 - 14/34

Author : Belovitskiy, G. Ye.; Romanova, T. A.; Sukhov, L. V.; and Frank,
I. M.

Title : Fission of uranium nuclei under the action of slow negative pi-
mesons and high-energy particles

Periodical : Zhur. eksp. i teor. fis. 28, 729-732, Jun 1955

Abstract : In this work the authors investigate the fission of uranium nuclei
by slow negative pi-mesons (G. Ye. Belovitskiy, et alii, Otechet
FIAN*, April 1950, June 1950, March 1951), by fast neutrons, with ener-
gies up to 460 Mev, and by gamma-rays with energies up to 250 Mev
(G. Ye. Belovitskiy et alii, ibid., Dec 1952). For the recording
of the fission of uranium nuclei they used photoplates with
emulsion layer 100 microns thick with uranyl acetate (T. A.
Romanova and G. Ye. Belovitskiy, ibid., June 1951), which plates
permitted the observation of protons with energies up to 30 Mev.
The irradiation of the plates by slow negative pi-mesons and fast
neutrons was carried out in the synchrocyclotron of the Institute
of Nuclear Problems. Academy of Sciences USSR; the irradiation by
gamma-rays was by the synchrotron of FIAN*. They note that the
energy spectrum of neutrons from "overcharging" (peresaryadka) of

Card 2/2

FD-2349

670-Mev protons on beryllium was measured by V. B. Flyagin. They present 5 photographs of indicated fission. They thank Prof. M. G. Meshcheryakov, G. P. Dzhalapov, and Ye. Grigor'yev for aid in experiments with negative pi-mesons and fast neutrons, and also thank Prof. V. I. Veksler and Yu. S. Ivanov for aid in experiments with gamma-rays of high energy. They state that a more detailed report on the results obtained will be published in this journal. They conclude that the distinguishing peculiarity of the process of fission of uranium nuclei at high energies of excitation is the significant probability of the emission of fast protons and alphaparticles; these particles bear only a comparatively small part of energy obtained by the uranium nucleus from the primary particle. Thirteen references.

Institution : Physical Institute imeni P. N. Lebedev, Acad. Sci. USSR (FIAN*)

Submitted : March 9, 1955

539.173
2385. FISSION OF URANIUM NUCLEI UNDER THE ACTION
OF SLOW π^- -MESONS, FAST NEUTRONS AND γ -RAYS OF
ENERGY UP TO 250 MeV. G.E. Belovitskii, T.A. Romasova,
L.V. Sukhov and I.M. Frank.
Zh. eksper. teoret. fiz., Vol. 29, No. 5(11), 537-50 (1955). In
Russian.

Events are studied in nuclear emulsions impregnated with uranium salt. The emulsions registered protons of energy less than 30 MeV. The plates were irradiated with π^- -mesons and neutrons of maximum energies 180 and 480 MeV, and with γ -rays of maximum energies 30, 80 and 250 MeV. In 96 fissions by π^- -mesons, 81 of the cases showed two fragments, 15 showed three. Probability of fission upon capture of a π^- -meson is estimated to be about 0.5. Similarly, fissions brought about by high-energy neutrons and γ -rays were analysed. Fission produced by all three of these high-energy agents is sometimes accompanied by a fast proton or α -particle. These, however, carry away only a small part of the energy released and, since the fission fragments have about the same energy distribution as those produced by slow neutrons, it is surmised that most of the energy is taken up in neutron emission.

G.E. Brown

Translation D 419421 - p.27

FRANK, I. M.

"Experimental Data on the Anisotropic Distribution of Fission Fragments",
a report presented at the Conference on the Physics of Nuclear Fission, 19-21
January 1956, Atom Energ., No. 1, 1956.

pmf

11

K. V.

[Signature]

Fission of the uranium nucleus under the influence of
slow γ mesons, fast neutrons and α rays with energies up
to 250 MeV

LSMEYANOV, A.N.; TOPCHIEV, A.V.; KURCHATOV, I.V.; SKOBEL'TSYN, D. .;
KAPITSA, P.B.; IOFFE, A.F.; VINOGRADOV, A.P.; ERENBURG, I.G.; TIKHONOV,
N.S.; FADYEYEV, A.A.; FRANK, I.M.; VEKSLER, V.I.; KORNEYCHUK, A.Ye.;
POPOVA, N.V.; LEHIDEVA, Z.A.; VASILEVSKAYA, V.L.; PETROVSKIY, I.G.;
ALEKSANDROV, A.D.; ARTSIMOVICH, L.A.; MESHCHERYAKOV, M.G.

Irene Joliet-Curie; obituary. Vest.AN SSSR 26 no.4:73-72 Ap '56.
(Joliet-Curie, Irene, 1897-1956) (MIRA 9:7)

USSR/ Physics

Card 2/2 Pub. 118 - 5/7

Author: Frank, I. M.

Title: Flash duration in the Vavilov-Chernokov effect

Periodical: Usp. Fiz. nauk, 58/1, 111-150, Jan 1956

Abstract: A special type of luminescence of pure liquids and some solid bodies, called the Vavilov-Chernokov effect, is considered. A short duration of the light flash plays the main role in the Vavilov-Chernokov effect. Various devices (counters) for measuring the light flash duration are described. A theory on which the counters have been constructed is presented. In accordance with this theory the duration of a light flash can be expressed as

Institution:

Submitted:

Card 2/2 Pub. 118 - 5/7

Periodical : Usp. Fiz. nauk, 58/1, 111-150, Jan 1956

Abstract : follows: $\tau_0 = \frac{1}{\Delta\nu}$, where the τ_0 is the time of the duration, and the $\Delta\nu$ is a narrow frequency band within which the Vavilov-Chernokov effect is observed. Nineteen references: 1 Eng., 1 Fr., 4 USA, 13 USSR (1924-1955). Diagrams; graphs.

65946

SOV/58-59-4-7684

24.6600

Translation from: Referativnyy Zhurnal Fizika, 1959, Nr 4, p 60 (USSR)

AUTHORS: Balabanov, Ye.M., Barit, I.Ya., Katsaurov, L.N., Frank, I.M., Shtranikh, I.V.

TITLE: Yield and Effective Cross-Section Measurements of $D(t,n)He^4$ and $D(d,p)T$ Reactions for a Thick Heavy-Ice Target

PERIODICAL: V sb.: Yadern. reaktsii na legkikh yadrakh. Moscow, Atomizdat, 1957, pp 48 - 56

ABSTRACT: The authors measured the yield and effective cross sections of $D(t,n)He^4$ and $D(d,p)T$ reactions for heavy ice in the 50 - 200 Kev deuteron energy range. A D_2^+ or HT^+ beam from an ion-accelerating tube was sorted in accordance with the different masses of the particles by means of a magnet and a system of diaphragms. The reaction yield was determined from the number of alpha-particles or protons registered at an angle of 90° to the beam with the aid of proportional counters. For the $D(t,n)He^4$ reaction a maximum was observed for 160 Kev tritons; the magnitude of the cross section at the maximum was equal to 4.34 barn. The yield and cross-section measurements of the $D(d,p)T$ reactions were carried out by way of a check,


Card 1/2

65946

SOV/58-59-4-7684

Yield and Effective Cross-Section Measurements of $D(t,n)He^4$ and $D(d,p)T$ Reactions for a Thick Heavy-Ice Target

since reliable results for this reaction using a gas target have been published (Sanders et al, Phys. Rev., 1950, Vol 77, p 1754, McNeill, K.G., et al, Phys. Rev., 1951, Vol 81, p 602). The results of the measurements showed that for a significant part of the energy range the obtained cross sections were 10 - 20% less than those obtained using a gas target. The authors assume that this is due to an inaccuracy in the values utilized for the energy losses in D_2O , or to some other systematic errors .

V.I.Ch. 

Card 2/2

FRANK, I.M.

Anisotropy in nuclear fission processes. Atom.energ.supplement
no.1:58-73 '57. (MIRA 10:10)
(Nuclear fission)

BALABANOV, Ye.M.; BARIT, I.Ye.; KATSAUROV, L.N.; FRANK, I.M.; SHTRANIKH, I.V.

Measurement of the effective cross section of the $D(t,n)He^4$ reaction
in the 40-730 Kev deuteron energy range. Atom. energ. suppl. no.5:57-
70 '57. (MIRA 11:2)

(Nuclear reactions) (Deuterons)

53-3-2/6

AUTHORS:

FRANK, I. M.
Skobel'tsyn, D.V., Frank, I.M.

TITLE:

The Physics Institute imeni P.N. Lebedev AN USSR (Fizicheskiy institut imeni P.N. Lebedeva)

PERIODICAL:

Uspekhi Fiz. Nauk, 1957, Vol. 63, Nr 3, pp. 503 - 525 (USSR)

ABSTRACT:

This institute dates back to the oldest center of experimental physical research in Russia. In 1725 the "Physical Cabinet" was founded together with the Academy of Science; in 1912 it was re-organized as physical laboratory. In 1921 this laboratory was transformed into the physical department of the physical-mathematical institute of the AN. In 1934 the institute, together with the AN, moved from Leningrad to Moscow, and in the same year the physical institute AN (FIAN) imeni P.N. Lebedev was established. The first director of the institute (1934-1951) was S.I. Vavilov, and after his death D.V. Skobel'tsyn. Special mention is made of Vavilov's works on the luminescence of uranyl salts which led to the discovery of the Vavilov-Cherenkov effect. The research work carried out and the collaborators of the various laboratories of this institute are mentioned. During World War II the institute was at Kazan, and by its activities assisted the armament industry.

Card 1/4

The Physics Institute imeni P.M. Lebedev AN USSR

53-3-2/6

The institute has now seven times the number of collaborators it had in 1945. At present 20 doctors and 100 candidates of sciences are working at the institute. The library at present contains 200 000 volumes (books and periodicals).

General survey of themes and of some of the results achieved by the institute: At first the success achieved by V.I. Veksler in connection with the construction of particle accelerators is mentioned. The successes achieved in the physics of elementary particles were prepared by the study of cosmic radiation. In recent times successful investigations were carried out by means of the photo-emulsion method. Also the theoretical department of the institute deals with the physics of elementary particles and of nuclear interactions. Well-known works are those by I.Ye. Tamm on mesonic interaction. Much is said about the work carried out by the institute in connection with the Vavilov-Cherenkov effect. The work and the themes dealt with by individual laboratories of the institute are then discussed.

Theoretical physics: Elementary particles, non-local theory, renormalizability and dispersion relations, multiple production of particles, cosmic rays, neutron physics, high-energy processes, solid bodies, low temperatures.

Card 2/4

The Physics Institute imeni P.N. Lebedev AN USSR

53-3-2/6

Nuclear Physics: Work concerning nuclear physics and cosmic rays has been developing since 1933, and supervision of this work was soon taken by D.V. Skobel'tsyn. During the war work on cosmic radiation was stopped until 1944, and already in 1944 the first expedition was sent to Pamir, and in 1947 the first high mountain station was established. The next report deals with new accelerators. Apart from theoretical work also theoretical research work was carried out on a large scale under supervision of M.A. Markov in the two following directions: Investigation of phenomena that are important for the research of atomic structure by the method of photonuclear reactions. Investigation of the properties of the meson field and of meson-producing processes by photonuclear interaction.

Further chapters deal with work carried out in the field of radiophysics, luminescence and optics, semiconductors and di-electrics. At present the effect produced by neutrons and fast electrons upon germanium is being studied. In this connection also some rules concerning the transformation of the energy of B-radiation into electric energy were set up. There are 67 references, all of which are Slavic.

Card 3/4

The Physics Institute imeni P.N. Lebedev AN USSR

53-3-2/6

AVAILABLE: Library of Congress

Card 4/4

DENISOV, F.P., red.; LAZAREVA, L.Ye., red.; LEYKIN, Ye.M., red.; ROZHANSKIY, I.D., red.; ~~FRANK, I.M., red.~~; SHAPIRO, I.S., red.; SHAPIRO, F.L., red.; POLENOVA, T.P., tekhn. red.

[Low and intermediate energy nuclear reactions; transactions of the conference] Yadernye reaktsii pri malykh i srednikh energiakh; trudy konferentsii. Moskva, Izd-vo Akad. nauk SSSR, 1958. 614 p.
(MIRA 11:12)

1. Vsesoyuznaya konferentsiya po yadernym reaktsiyam pri malykh i srednikh energiakh. Moscow, 1957.
(Nuclear reactions)

FRANK, I. M.

"The Application of Atomic Energy for Peaceful Purposes"

Lecture to be delivered by Soviet Scientists at the Brussels Exhibition, August 1958. The delivered lectures will be available in English, French, Flemish and German as individual brochures.
(Priroda, 1958, No. 8, p. 116)

24(5)

PHASE I BOOK EXPLOITATION

SOV/2121

Frank, Il'ya Mikhaylovich, Corresponding Member, USSR Academy of Sciences, Laureate of the Nobel Prize

Izlucheniye Vavilova-Cherenkova; lektsiya, pročitannaya na Vsemirnoy vystavke v Bryussele... (Vav'lov-Cherenkov Radiation; a Lecture Delivered at the Brussels' World Fair on August 13, 1958) Moscow, Izd-vo "Znaniy", 1959. 30 p. (Series: Vsesoyuznoye obshchestvo po rasprostraneniyu politicheskikh i nauchnykh znaniy. Seriya IX, 1959, no. 8) 36,000 copies printed.

Sponsoring Agency: Vsesoyuznoye obshchestvo po rasprostraneniyu politicheskikh i nauchnykh znaniy.

Ed.: I.B. Faynboym; Tech. Ed.: L.Ye. Atroshchenko.

PURPOSE: The booklet is intended for the general reader interested in physics.

Card 1/2

Vavilov-Cherenkov Radiation (Cont.)

SOV/2121

COVERAGE: This booklet contains a lecture on luminescence phenomenon delivered by the author at the Brussels' World Fair on August 13, 1958. The author describes the Vavilov-Cherenkov effect and the properties of fast particles at some length. Brief biographies of Soviet scientists who have been awarded Nobel prizes are included in the booklet. No references are given.

TABLE OF CONTENTS:

On Light and Matter	3
Luminescence	4
Vavilov-Cherenkov Effect	6
Nature of the Phenomenon	9
Vavilov-Cherenkov Radiation and the Properties of Fast Particles	12
Nobel Prizes Awarded to Soviet Scientists (information)	20

AVAILABLE: Library of Congress

Card 2/2

TM/ad
8-21-59

FRANK, I. M.

21(0), 22(0) PHASE I BOOK EXPLOITATION 307, 32
Akademiya nauk SSSR. Fizicheskii institut
Issledovaniya po eksperimental'noy i teoreticheskoy fizike; (bornil-
(Studies on Experimental and Theoretical Physics; Collection of
Articles) Moscow, Izd-vo AN SSSR, 1959. 304 p. Errata slip
inserted. 2,300 copies printed.

Ed.: I. L. Fabelinskii, Doctor of Physical and Mathematical Sci-
ences; Eds. of Publishing House: A. L. Chernyak and V. G. Bergeyau.
Tech. Ed.: Yu. V. Rykina; Commission for Publishing the Collection
in Memory of Grigoriya Samoilovich Lur'e, Institute of Physics, USSR
Academy of Sciences.
S. A. Mandel'shtam, Doctor of Physical and Mathematical Sciences;
S. I. Mandel'shtam, Doctor of Physical and Mathematical Sciences;
I. L. Fabelinskii, Doctor of Physical and Mathematical Sciences;
P. S. Landsberg-Baryshanskaya, Candidate of Physical and Math-
ematical Sciences; and G. P. Motulevich (Secretary), Candidate of
Physical and Mathematical Sciences.

PURPOSE: This book is intended for physicists and researchers
engaged in the study of electromagnetic radiations and their role
in investigating the structure and composition of materials.
COVERAGE: The collection contains 30 articles which review
investigations in spectroscopy, optics, molecular optics, semi-
conductor physics, nuclear physics, and other branches of
physics. The introductory chapter gives a general profile
of G. S. Landsberg, Professor of the Department of
Optics of the Division of Physical Technology at Moscow Uni-
versity, and reviews his work in Rayleigh scattering, combat
gases, spectral analysis of metals, etc. No personalities are
mentioned. References accompany each article.

Report on B. S. Kinetics of the Action of Light Gases on the Intensity of Absorption Spectra of Vapors of Aromatic Com- pounds	149
Obrazov. I. V. and Ye. S. Trukhor. The Resistance of Mica to Rupture Along the Cleavage Plane	159
Rytov, S. M. The Correlation Theory of Rayleigh Light Scat- tering	175
Sobel'man, I. I. The Quantum Mechanics Theory of the Intensity of Combined-Scattering Lines	192
Sushchinskii, M. M. Dependency of the Width of Combined- Scattering Lines of the Anisotropy of a Derived Polarizability Tensor	211
Tam, I. Ye. Present State of the Theory of Weak Interactions of Elementary Particles	218
Turman, L. A. and B. A. Chaynov. The Illumination of Dielectrics in High Voltage a-c Electric Fields	231
Ukhollin, S. A. and M. Z. Promina. Investigation of Combined Light-Scattering Spectra in H ₂ O-H ₂ O and H ₂ O ₂ -Dioxine Solutions	244
Fabelinskii, I. L. The Thin Structure of Lines of Rayleigh Light-Scattering in Gases	254
Frank, I. M. The Role of the Group Speed of Light in Irradia- tion in a Refractive Medium	261
Frieh, S. E. and I. P. Bogdanova. Excitation of Spectral Lines in the Negative Illumination of a Gas Discharge	275
Frisberg, A. A. and V. Y. Medler. The Possibility of Increasing the Sensitivity of the Spectral Determination of Some Elements	287
Shpol'skiy, E. Z. The Interpretation of Spectra of Aromatic Hydrocarbons in Frozen Crystalline Solutions	296

NUMBER: 31(10)
AUTHOR: Parthiv'ko, V.
TITLE: The Fifth Session of the Scientific Council of the Joint Institute of Nuclear Research (Pratsya sessiya Dvenadtsati sosednykh nauchnykh uchebnykh institutov)
DATE: 1967-09-04-16/27

PERIODICAL: Atomnaya energiya, 1959, Vol. 6, No. 4, p. 479 (1958).

ABSTRACT: The fifth session of the Scientific Council of the Ob'yedinsenny Institute yadernykh issledovaniy (Joint Institute of Nuclear Research) was held from January 4 to 17, 1959. Lectures were held on the following important papers, which were also discussed: Professor V. Babelov spoke about neutrino yadernykh obtained by the work carried out by the author; investigations of the mechanism of induced isobaric scattering of neutrons were carried out of the fast neutrons; scattering of mesons on polarized and unpolarized particles, scattering of mesons on polarized mesons, processes of weak interaction in the decay of π -mesons, and the properties of π -mesons. The Director of the Laboratory, V. I. Veksler, gave a report on the work of the Laboratory for Theoretical Physics, Akademskiy N. S.

Card 1/3

Card 3/3

24(4)

AUTHOR:

Frank, I. M.

SOV/56-36-3-25/71

TITLE:

On the Role of the Group Velocity of Light in the Case of Radiation in a Refractive Medium (O roli gruppovoy skorosti sveta pri izluchении v prelomlyayushchey srede)

PERIODICAL:

Zhurnal eksperimental'noy i teoreticheskoy fiziki, 1959, Vol 36, Nr 3, pp 823-831 (USSR)

ABSTRACT:

In the present paper the author investigates the part played by the group velocity of light in a radiation, the source of which moves uniformly in a refractive isotropic medium, it being assumed that the medium is transmissive with respect to the emitted light. The radiation of a moving light source depends essentially on the ratio between the velocity of the emitter and the phase velocity of the emitted frequency. For particles which generate a time-constant electromagnetic field (e.g. electric charges or constant magnetic dipoles), i.e. at an eigenfrequency that is equal to zero, the Cherenkov effect occurs (which is also called Vavilov-Cherenkov effect in Russian publications), if the velocity of motion v is equal to or begins to exceed the phase velocity of light. In the case of

Card 1/2

On the Role of the Group Velocity of Light
in the Case of Radiation in a Refractive Medium

SOV/56-36-3-25/71

this connection between the direction of light radiation and the emitted frequency ω_0 it holds that $v \cos \theta / (c/n(\omega_0)) = 1$, $n(\omega_0) = c/v \cos \theta$, $u = c/n(\omega_0)$;

the emitted frequency, like Cherenkov radiation, is subjected to a Doppler effect for which $v/(c/n(\omega_0)) = \beta n(\omega_0)$ holds.

Basing on these assumptions the author first investigates the Cherenkov effect and the conditions for the occurrence of complex radiation. In the last part of the paper the connection between the velocity of the motion of the emitter and the group velocity of light is investigated, which leads the author to the conclusion that the emitter is not able to catch up with the light signal emitted by it in the direction of its motion. There are 3 figures and 4 Soviet references.

ASSOCIATION: Fizicheskiy institut im. P. N. Lebedeva Akademii nauk SSSR
(Physics Institute imeni P. N. Lebedev of the Academy of Sciences, USSR)

SUBMITTED: August 16, 1958

Card 2/2

24 (4)

AUTHOR:

Frank, I. M.

SOV/53-68-3-4/11

TITLE:

The Optics of Light Sources Moving in Refracting Media (Optika
istochnikov sveta, dvizhushchikhsya v prelomlyayushchikh sredakh)

PERIODICAL:

Uspekhi fizicheskikh nauk, 1959, Vol 68, Nr 3, pp 397-415 (USSR)

ABSTRACT:

This is a reproduction of a lecture delivered on the occasion of the awarding of the Nobel Prize. It consists of five parts. In part 1 the particular features of the radiation discovered by Cherenkov and Vavilov in a refracting medium are discussed together with the conditions for the occurrence of this radiation and its properties. In this connection reference is made to L. I. Mandelstam and V. L. Ginzburg. In part 2 the author discusses as a characteristic example the so-called transition radiation; it occurs e.g. if a uniformly moving charged particle exceeds the boundary of two media with different refraction indices. The intensity of this radiation is proportional to the kinetic energy of the particle at low velocities, and at relativistic velocities it grows like the logarithm of the total energy. Two particularities of this radiation within the range of ultrarelativistic velocities are discussed and the results obtained by investigations carried out by A. Ye. Chudakov and

Card 1/4

The Optics of Light Sources Moving in Refracting Media SOV/53-68-3-4/11

V. Ye. Pafomov are mentioned. Part 3 deals with the radiation spectrum and with the quantum-theoretical interpretation of the Cherenkov effect (according to Ginzburg). A number of the most important equations is given. Again, a particle moving with constant velocity in a refracting medium is investigated. If the momentum $n\hbar\omega/c$ of the photon is small compared to the momentum of the radiation source, the theorem of the conservation of momentum may be written down as

$\frac{n\hbar\omega}{c} \cos \theta = \frac{\Delta E}{v}$, where ΔE denotes the kinetic energy of the radiation source. $\Delta E = \hbar\omega \pm \hbar\omega_0$ (ω_0 is the eigenfrequency in the laboratory system). Thus, $\frac{n\omega}{c} \cos \theta = \frac{\omega \pm \omega_0}{v}$ is obtained. In the following, 3 cases are discussed: if $\frac{n\omega}{c} \cos \theta = 1$,

< 1 and > 1 . Finally a classical discussion of the Doppler effect is given, and the possibility of a velocity greater than that of light in a medium is discussed. It was found that an arbitrary system capable of interacting with radiation slows down at a velocity greater than that of light and emits light. In part 4 the author discusses radiation thresholds and investigates the conditions for the occurrence of Cherenkov

Card 2/4

The Optics of Light Sources Moving in Refracting Media SOV/53-68-3-4/11

radiation and of the Doppler effect for velocities greater than that of light. For this, the condition $\frac{vn(\omega)}{c} > 1$ is, first of all, given, which says that v must be greater than the phase velocity of light. L. I. Mandel'shtam pointed out that the complex Doppler effect is connected with the amount of the group velocity W of light. On the basis of a diagram (Fig 1), which represents the frequency dependence of the wave vector $k(\omega) = \frac{\omega n(\omega)}{c}$, these conditions are discussed and a number of formulas is derived. Figure 2 shows some possible cases of connections existing between \vec{u} , \vec{v} , \vec{w} and θ , which are discussed. Part 5 finally deals with Cherenkov radiation in optically anisotropic media. In this connection the case is investigated in which the radiation source moves in a crystal, in which the refraction index depends not only on light frequency but also on the angle and on polarization. The influence exercised by the latter is discussed in detail, and a number of special cases is further discussed on the basis of schematical drawings. Also the Doppler effect is dealt with, and attention is drawn

Card 3/4

The Optics of Light Sources Moving in Refracting Media SOV/53-68-3-4/11

to the most important investigations carried out by K. A. Barsukov and A. A. Kolomenskiy. There are 6 figures and 18 references, 16 of which are Soviet.

Card 4/4

PHASE I BOOK EXPLOITATION SOV/4393

Cherenkov, Pavel Alekseyevich, Professor, Igor' Yevgen'yevich Tamm, Academician, and Il'ya Mikhaylovich Frank, Corresponding Member, Academy of Sciences USSR

Nobelevskiye lektsii (Nobel Prize Papers) Moscow, Fizmatgiz, 1960. 73 p. 7,000 copies printed.

Ed.: T. V. Mikhalkovich; Tech. Ed.: Ye. A. Yermakova.

PURPOSE: This pamphlet is intended for physicists and researchers engaged in the application of the Cherenkov radiation principle in experimental physics.

COVERAGE: The pamphlet contains lectures by Professor P. A. Cherenkov, Academician I. Ye. Tamm, and Corresponding Member of the USSR Academy of Sciences I. M. Frank given in Stockholm on December 11, 1958 when receiving the Nobel Prize in physics. The supplementary article relates the history of the discovery of the Cherenkov radiation and presents biographical data on the three Nobel Prize re-

Card 1/3

Nobel Prize Papers

SOV/4393

ciipients. Photographs of the prize winners are included in the booklet. The complete text of the speeches and of the article were previously published in Uspekhi fizicheskikh nauk, v. 67, no. 1, and v. 68, no. 3. The articles are accompanied by bibliographies listing Soviet and other technical literature.

TABLE OF CONTENTS:

Editor's Preface

3

Cherenkov, P. A. Radiation of Particles Having Velocity Greater Than That of Light and Some Applications of This Radiation in Experimental Physics

5

Tamm, I. Ye. General Properties of Radiation Emitted by Systems Moving at Greater Velocities Than That of Light and Some Applications to the Physics of Plasma

20

Card 2/3

Nobel Prize Papers

SOV/4393

Frank, I. M. Optics of Light Sources Moving in Refractive Media

35

Supplement. Bolotovskiy, B. M. Soviet Scientists, Winners of the 1958 Nobel Prize in Physics

64

AVAILABLE: Library of Congress

Card 3/3

JA/rn/ec
10-20-60

H/016/60/010/010/002/004
B009/B057

AUTHOR: Frank, I. M.
TITLE: Optics of Light Sources Moving in Refractive Media
PERIODICAL: Fizikai Szemle, 1960, Vol. 10, No. 10, pp. 298-304

TEXT: This is an abridged text of a lecture delivered at the conferment of the Nobel Prize upon the author in Stockholm on December 11, 1958. The full-length text was published in Uspekhi Fizicheskikh Nauk, Vol. 68 (1959), pp. 397-415. 1) Characteristics of light in a medium: In vacuum, the velocity of light is always constant and higher than the velocity of the light source. In a refractive medium, the ratio of the velocity of the radiant to the velocity of wave propagation depends very much on the velocity of light in the medium and on its changes. As a result, not only the characteristics of radiation but sometimes also its appearance depends on the features of light propagation in the medium. In this paper, the simplest case of radiation in refractive media is considered, where the translational motion of the system may be supposed to be uniform and rectilinear. 2) Transition radiation: The theory of transition radiation

Card 1/3

Optics of Light Sources Moving in Refractive Media

H/016/60/010/010/002/004
B009/B057

was developed by V. L. Ginzburg and the author of this paper more than 10 years ago, but was not examined experimentally. Most promising results have been recently reached by A. Ye. Chudakov but have not been published as yet. V. Ye. Pafomov pointed out that for very small thicknesses of the radiator, transition radiation must be considered as a component in the observation of the Cherenkov radiation. 3) Spectrum and quantum-mechanical interpretation: The quantum-mechanical theory of transition radiation was first developed by Ginzburg who demonstrated that the classical formula of the cosine of the angle at which the radiation takes place may be supposed to be exact with a very small correction. In vacuum, motion at a velocity exceeding that of light is impossible. In a medium, it appears possible, but nature does not entirely cancel that prohibition. An arbitrary system, capable of radiation interactions, will - at velocities exceeding that of light - be slowed down by emission of light. 4) Thresholds of radiation: The spectrum of radiation is determined by the velocity of the system, its fundamental frequency, and the phase velocity of light valid for the system. From formulas deduced for the Cherenkov phenomenon and the Doppler effect at velocities exceeding that of light, it follows that these radiations can develop only if the

Card 2/3

Optics of Light Sources Moving in Refractive Media

H/016/60/010/010/002/004
B009/B057

velocity of motion of the system exceeds the phase velocity of light. This statement which holds for isotropic media determines the threshold of generation of radiation. In a refractive medium, energy transfer takes place not at phase but at group velocity. If the conditions of its development are satisfied, radiation develops and draws energy from the radiant. When the velocity of light reaches the threshold, radiation ceases and so does the discharge of energy. This will be the case when the group velocity of light equals the velocity of motion of the system. The role of the group velocity of light will manifest itself particularly clearly in an anisotropic medium where it includes a certain angle with the phase velocity. (Section 5 of the lecture dealt with radiation in optically anisotropic media and is excluded from this text). L. I. Mandel'shtam is mentioned. There are 12 references: 11 Soviet and 1 US.

Card 3/3

81670

S/056/60/038/06/05/012
B006/B056

24.3200

AUTHOR: Frank, I. M.

TITLE: Critical Velocity of Light Emission in Optically Anisotropic Media

PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki, 1960,
Vol. 38, No. 6, pp. 1751-1757

TEXT: The author deals with the investigation of the emission of a light source of arbitrary natural frequency, moving uniformly in an optically anisotropic transparent medium. In an optically isotropic medium a Vavilov-Cherenkov radiation of the frequency ω occurs, if the velocity of motion v of the emitting charge attains the phase velocity of light $u(\omega) = c/n(\omega)$. In the expression for the wave vector $k(\omega)$ only the phase velocity occurs; it is, however, to be assumed that in optically anisotropic medium the group velocity plays a similar part. First, the peculiarities occurring in an anisotropic medium are discussed, which are based mainly upon the fact that the phase velocity depends both on the polarization of the waves and on the direction of $\vec{k}(\omega)$. On

Card 1/3

81670

Critical Velocity of Light Emission in
Optically Anisotropic Media

S/056/60/038/06/05/012
B006/B056

the basis of the simple case of a monochromatic wave of given polarization it is shown that for the critical velocity $\vec{u}'(\omega)$ occurring in the case of an anisotropic medium, $\vec{v} = \vec{u}'$ holds. This second phase velocity u' is described by the author as "radiation velocity along the beam". Fig. 1

illustrates the foregoing. It holds generally that $\vec{k}'(\omega) = \frac{\omega}{u'^2} \vec{u}' = \frac{\omega \pm \omega_0}{v^2} \vec{v}$;

if $\omega_0 = 0$, $\vec{v} = \vec{u}'$ is thus satisfied. This equation is the condition

holding in an anisotropic medium for the occurrence of a radiation of the frequency ω . In the case represented in Fig. 1, the surface of the wave vectors is an ellipsoid, and therefore the "extraordinary" beam in a uniaxial crystal is concerned. An analogous investigation is carried out for the ordinary beam (k - surface: sphere); $v = u = u'$. In the following, the peculiarities occurring in a biaxial crystal are discussed. Fig. 2 shows k' as a function of ω for a given polarization. If w is the group velocity of a beam of the frequency ω , which runs the direction \vec{v} , the condition holding for the occurrence of a radiation or of new components of it is $\vec{v} = \vec{w}(\omega_{gr})$, where ω_{gr} is the frequency

Card 2/3

81670

Critical Velocity of Light Emission in
Optically Anisotropic Media

S/056/60/038/06/05/012
B006/B056

primarily occurring in the spectrum, for which the condition given above for $k'(\omega)$ is satisfied. Finally, the case $\omega < \omega_0$ is investigated. All results obtained are analogous to those obtained for an isotropic medium (Refs. 5, 6). They hold for the case in which the direction of motion coincides with the direction of the beam. V. L. Ginzburg and V. Ye. Pafomov are mentioned in this paper. There are 2 figures and 6 Soviet references.

ASSOCIATION: Fizicheskiy institut im. P. N. Lebedeva Akademii nauk SSSR
(Physics Institute imeni P. N. Lebedev of the Academy of
Sciences USSR)

SUBMITTED: December 21, 1959

Card 3/3

✓

FRANK, I. M., SPALING, F. L., KAZVITSKIY, Y. G., BLOKHIN, D. I.,
BLOKHIN, G. P., LYUMFINA, Y. A., BONDARENKO, I. I., BELYACHIN, P. V.,
ZAINOVSKIY, A. S., ZINGOV'YEV, V. P., KAZACHOVSKIY, O. B., KRAZNOYAROV, N. V.,
LEYPUNSKAYA, A. I., MALIK, V. A., MAZAROV, P. M., MECLAYEV, S. K.,
SEVILESKIY, Y. Y., UKRAINSKIY, F. I.

"A Pulse fast reactor."

report submitted for the IAEA seminar on the Physics of Fast and
Intermediate Reactors, Vienna, 3-11 Aug 1961.

Acad Sci. USSR Moscow

21.1910 21.4210
26.2200
AUTHORS: Blokhin, G. Ye., Blokhintsev, D. I., Blyumkina, Yu. A.,
Bondarenko, I. I., Deryagin, B. N., Zaymovskiy, A. S.,
Zinov'yev, V. P., Kazachkovskiy, O. D., Kim Khen Bon,
Krasnoyarov, M. V., Leypunskiy, A. I., Malykh, V. A.,
Nazarov, P. M., Nikolayev, S. K., Stavisskiy, V. Ya.,
Ukrainsev, F. I., Frank, I. M., Shapiro, F. L.,
Yazvitskiy, Yu. S.

22873
S/089/61/010/005/001/015
B102/B214

TITLE: A pulsed fast reactor

PERIODICAL: Atomnaya energiya, v. 10, no. 5, 1961, 437-446

TEXT: The present paper gives a description of the pulsed fast reactor of the Ob'yedinennyi institut yadernykh issledovaniy (Joint Institute of Nuclear Research) which became critical in June, 1960. This reactor, called W6P (IBR) reactor, serves as pulsed fast neutron source (mean power ≈ 1 kw) for physical investigations, particularly for time-of-flight experiments. Its most distinguishing feature is the very small contribution ($\sim 10^{-4}$) of the delayed neutrons in its normal operation; it is about

Card 1/7 ✓

22873

S/089/61/010/005/001/015
B102/B214

A pulsed fast reactor

one hundredth of that of the usual steady uranium reactor. The pulses appear because whenever the reactor becomes overcritical a burst of prompt neutrons results. The half width of these pulses is 36 μ sec. The frequency with which the pulses are repeated can be varied between 8 and 80 pulses/sec. Fig. 2 shows the construction of this reactor. The periodic change in the reactivity is brought about by the displacement of the two U^{235} blocks placed in two disks that can be rotated. The main block is pressed in the form of a disk, 1100 mm in diameter, and can be rotated with a peripheral velocity of 276 m/sec (at 6000 rpm) during which it passes through the core center. The reactivity change obtainable from the motion of the main block is 7.4 %, that obtainable from the motion of the auxiliary block is 0.4 %. The stationary part of the core consists of plutonium lumps in steel jackets. The reactor is started by a rough regulator, in this case a movable part of the reflector. It gives a reactivity change at the rate of $13 \cdot 10^{-5} - 1.3 \cdot 10^{-5} \text{ sec}^{-1}$. The manually operated rod is also a part of the reflector. Two plutonium rods in electromagnetic suspension serve as scram. They can be separated from the core with an acceleration of 20 g. Their separation causes a reactivity

Card 2/24

22R73

3/009/61/010/005/001/015
B102/B214

A pulsed fast reactor

decrease of 2-1.1 %; the rough regulator allows a reactivity change of 2.4 %, the manual regulator 0.1 %, and the automatic regulator 0.036 %. The reactor possesses also a reactivity booster for the production of one intensive pulse. The control and shield system is an automatically functioning electronic arrangement with BF_3 counters and ionization chambers. The whole reactor is placed in a room of size 10-10-7 m whose concrete walls allow complete protection from radiation. The most important experimental arrangement consists of a 1000 m long neutron conductor, a metal tube, 400 mm in diameter in the first part and 800 mm in the second part in which a pressure of 0.1 mm Hg is maintained. This conductor connects a chain of so-called "intermediate pavilions" (at distances of 70, 250, 500, 750, and 1000 m from the reactor) in which experiments can be carried out. There is also an additional neutron conductor of 100 m length. The reactor chamber is joined to an experimental chamber in which four neutron beams of up to 800 mm diameter are available. There is such an experimental chamber also above the reactor chamber. Various experiments were carried out with the reactor and they are described in the present paper. These are experiments with stand

Card 3/74

A pulsed fast reactor

22873
S/089/61/010/005/001/015
B102/B214

assemblies and slowly moving main block for the determination of the most important parameters of the reactor; experiments with a core assembly (unmoved), experiments with rotating (5000 rpm) main block and a Ra- α -Be source in the core for the investigation of the effect of the multiplication factor, etc. The most important results are represented graphically. For example, Fig. 8 shows the dependence of the half width Θ of a pulse on the reactivity; the dashed line holds for the quasistationary case, the dot-dash line for the case of $\Theta = K(\tau/a)^{1/3}v^{-2/3}$, where v is the velocity of motion of the (rotating) main block; in the quasistationary case $\Theta = 2\sqrt{\epsilon_m/av^2}$, where ϵ_m is the reactivity at the maximal multiplication factor; $\epsilon = \epsilon_m - ax^2$, where x is the displacement of the main block. The reactor has been actually used for the measurement of the total, scattering, capture, and fission cross sections by the time-of-flight method. Further experiments will be carried out with a view to obtaining increase of power and decrease of the pulse duration. There are 15 figures and 3 references: 2 Soviet-bloc and 1 non-Soviet-bloc. The reference to the English-language publication reads as follows: J. Orndorf, Nucl. Sci. and Engng, 2, No. 4, 450 (1957).

Card 4/14

LEVSHIN, V.L.; TEREVIN, A.N.; FRANK, I.M.

Progress of S.I.Vavilov's work in the field of physics.

Usp.fiz.nauk 75 no.2:215-225 O '61.

(MIRA 14:10)

(Vavilov, Sergei Ivanovich, 1891-) (Physics)

FRANK, I.M.

Transient radiation and the Vavilov-Cherenkov effect. Usp.fiz.nauk
75 no.2:231-240 0 '61. (MIRA 14:10)
(Cherenkov radiation)

GRIGOROV, Naum Leonidovich; KONDRAT'YEVA, Marina Aleksandrovna;
RAPOPORT, Ilya Davidovich; FRANK, I.M., red.; GRIGOROVA,
V.A., red.; FLAKSHE, L.Yu., tekhn. red.

[Cosmic rays]. Kosmicheskie luchy. Moskva, Fizmatgiz. 1962.
83 p. (Praktikum po iadernoi fiziki, no.2).

(MIRA 1617)

1. Chlen-korrespondent AN SSSR (for Frank).
(Cosmic rays)

ESTULIN, Isay Veniaminovich; ZHABOTINSKIY, Ye.Ye., red.; FRANK, I.M., red.;
MURASHOVA, N.Ya., tekhn. red.

[Radioactive radiations]. Radiativnye izlucheniia. Moskva,
Fizmatgiz, 1962. 260 p. (Praktikum po iadernoi fizike, no.1).
(MIRA 16:4)

1. Chlen-korrespondent AN SSSR (for Frank).
(Radioactivity)

FRANK, I.M., otv. red.; DAVYDOV, A.S., red.; LAZAREVA, L.Ye., red.
NEMIROVSKIY, P.E., red.; CHUYEV, V.I., red.; POLYAKOVA, T.V.,
tekhn. red.

[Transactions of the Second All-Union Conference on Nuclear
Reactions at Low and Medium Energies] Trudy Vtoroy Vsesoyuznoy
konferentsii po yadernym reaktsiyam pri mal'kikh i srednikh ener-
giyakh, Moscow, 1960. Moskva, Izd-vo Akad. nauk SSSR, 1962.
658 p. (MIRA 16:2)

1. Vsesoyuznaya konferentsiya po yadernym reaktsiyam pri ma-
lykh i srednikh energiyakh, 2d, Moscow, 1960.
(Nuclear physics—Congresses)

S/903/62/000/000/014/044
B102/B234

AUTHORS: Benetskiy, B. A., Betin, Yu. P., Bukarev, V. A., Frank, I. M.

TITLE: (n, γ) -correlation in inelastic scattering of 14-Mev neutrons from C^{12} nuclei

SOURCE: Yadernyye reaktsii pri malykh i srednikh energiakh; trudy Vtoroy Vsesoyuznoy konferentsii, iyul' 1960.g. Ed. by A. S. Davydov and others. Moscow, Izd-vo AN SSSR, 1962, 178-179

TEXT: 14-Mev neutrons from a $T^3(d,n)He^4$ source were scattered from a toroidal carbon scatterer; the γ -rays were recorded by a NaI(Tl) scintillation detector with $\Phi BY-29$ (FEU-29) photomultiplier, the neutrons by a stack of plates of an organic scintillator separated by plexiglas and connected with an $\Phi BY-24$ (FEU-24) multiplier. Also the recoil protons with $7 \text{ Mev} \leq E_p \leq 14 \text{ Mev}$ were recorded. The pulse-height resolution of the γ -detector was 10% for Tn^{65} 1.12-Mev quanta; the coincidence circuit had a time resolution of $2 \cdot 10^{-7}$ sec. The γ -spectrum was analyzed with the help of a pulse-height analyzer. The angular distribution of the 4.4-Mev quanta emitted on the transition of the C^{12} nucleus from the first excited to the

Card 1/2

(n, γ)-correlation in inelastic...

S/903/62/000/000/014/044
B102/B234

Ground state ($2^+ \rightarrow 0^+$) could be described by $f(\lambda) = A + \sin^2(\lambda - \lambda_0)$ (cf. Ann. Phys., 2, 471, 1957) with $A = (6.27 \pm 0.14)$ and $\lambda_0 = (80 \pm 13)$. The anisotropic part of the $f(\lambda)$ function has the same character, independent whether the angle of emission of inelastically scattered neutrons is fixed or not.

ASSOCIATION: Fizicheskii institut im. P. N. Lebedeva AN SSSR (Physics
Institute imeni P. N. Lebedev AS USSR)

Card 2/2

L 16138-63 EWT(m)/BDS AFFTC/ASD
ACCESSION NR: AT3001852

S/2504/62/014/000/0117/0146

AUTHOR: Frank, I. M.

TITLE: A pulse method for the investigation of the properties of slow neutrons 19

SOURCE: AN SSSR. Fizicheskii institut. Trudy, v.14, 1962, 117-146

TOPIC TAGS: neutron, slow, pulse, flashing, spectroscopy, prism, fast, flight time, life time, diffusion coefficient, attenuation, moderation, moderated

ABSTRACT: This classical theoretical paper discusses the possible application of a high-voltage ion tube operating in a pulsed regime for the study of the properties of slow neutrons. The paper reproduces a report of the Fizicheskii institut AN SSSR (Physics Institute, AN SSSR) dated March 1954. The material of this study, together with the results of an experimental investigation on the same subject, was utilized in the report of A.V. Antonov, A.I. Isakov, I.D. Murin, B.A. Neupokoyev, I.M. Frank, F.L. Shapiro, and I.V. Stranikh at the First Geneva Conference on the Peaceful Use of Atomic Energy. The paper is reprinted essentially without any alterations. The pulse method can be employed most naturally for the determination of the maximum magnitude of the diffusion coefficient D of neutrons and its dependence on the conditions of the experiment. Inasmuch as in many instances not only D but also the life time of the neutron, T , can be determined, $L^2 = DT$ can thus

Card 1/4

L 16138-63

ACCESSION NR: AT3001852

also be obtained. The method provides indications on the time required for the establishment of a stationary velocity spectrum of neutrons and on the behavior of various groups of slow neutrons. More complete information on the properties of a neutron gas can be obtained by the method of the "pulsed prism" which permits the determination of the moderation (deceleration) parameter τ . Section 1 provides a survey of existing neutron-moderator theory, especially in the light of the effect of a pulsed emission regime. Sec. 2: The application of a pulse-type neutron source for the study of the diffusion of a neutron gas. Sec. 3: The initial distribution of decelerating neutrons. Sec. 4: The density attenuation of slow neutrons. Sec. 5: A pulse-type neutron source in a prism. Sec. 6: Application of two-group theory in the case of a pulse-type neutron source. Sec. 7: Conclusions: (1) The principal application of a pulse-type neutron source at the present time appears to be in flight-time neutron spectrometry. The use of $T(d,n)^4\text{He}$ in the "flashing-tube" method opens broad perspectives in various areas of neutron physics. Possible applications: Assuming that fast neutrons from a pulse-type source enter a system comprising a test substance and having a prescribed shape, for example a cube, a parallelepipedon, or a sphere, and dimensions that differ not too greatly from the neutron moderation length, the exponential law of the density attenuation of the neutron gas, beginning at a certain time point, permits a direct determination of the diffusion coefficient D (assuming the life time T to be

Card 2/4

L 16138-63

ACCESSION NR: AT3001852

approximately known). (2) In substances in which the velocity distribution of moderated neutrons becomes stationary within a time that is less than the lifetime of the slow neutrons, the magnitude of D will also attain a constant value. Inasmuch as in such substances D is independent of Ω , that is, the dimensions of the system, the magnitude of D becomes a physical constant that characterizes the neutron properties of the substances investigated. The magnitude of the attenuation coefficient, α , will then depend linearly on Ω ; the slope of the α -vs.- Ω line obtained by means of measurements in systems of various sizes yields D , while its initial ordinate, obtained by extension of the line to $D=0$, provides the magnitude of $1/T$. This method requires considerably less test material than the customarily employed prism method. (3) In extremely small systems, that is, in systems having a large Ω , it may be anticipated that the magnitude of D_{eff} , as determined from the magnitude of α , will decrease with an increase in the dimensions of the system. This peculiarity can be employed for the clarification of the properties of the various groups of slow neutrons. (4) The exponential behavior of the density of the neutron gas in the system is not established at once; the investigation of the duration of the transition stage can provide indications relative to the duration of the establishment of a stationary velocity distribution of the decelerating neutrons. (5) The most complete indication on the moderation (deceleration) and the diffusion of neutrons can be obtained by the method of the "flashing"

Card 3/4

L 16138-63

ACCESSION NR: AT3001852

4
prism, that is, by placing a pulse-type source inside a prism. (6) In some instances, for example, for water, it may be useful to employ a system consisting of a plate of prescribed thickness and great area (approximating a two-dimensionally infinite system). "In the course of the performance of this study, and in the discussion of its results, many valuable observations were made by F. L. Shapiro, to whom I express my warmest gratitude. I also thank M. V. Kazar-novskiy and A. V. Antonov for a number of remarks made in the discussion of the work." Orig. art. contains 2 figs., 2 tables, and 96 numbered equations.

ASSOCIATION: Fizicheskii institut AN SSSR (Physics Institute, AN SSSR)

SUBMITTED: 00	DATE ACQ: 11Apr63	ENCL: 00
SUB CODE: PH	NO REF SOV: 007	OTHER: 007

Card 4/4

S/056/63/044/002/009/065
B102/B186

AUTHORS: Benetskiy, B. A., Frank, I. M.

TITLE: Investigation of the angular correlation between γ -photons and 14-Mev neutrons inelastically scattered from carbon nuclei

PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki, v. 44, no. 2, 1963, 454-461

TEXT: The possibilities of $n'\gamma$ angular correlations are discussed in order to obtain an unambiguous answer to the question as to whether the process $C^{12}(n,n'\gamma)C^{12}$ occurs via compound nucleus formation or via direct interaction. It is found that in the latter case a state of definite parity (2^+) will result and the distribution will be symmetrical with respect to $(\psi_n - \pi)/2$, i.e. the correlation will be characterized by $\sin^2(\theta_\gamma - \psi_n/2 - n\pi/4)$. The $n'\gamma$ correlation was measured in a simple arrangement: the neutrons were obtained from a DT source ($E_n = 14.2$ Mev);

Card 1/19

Investigation of the angular ...

S/056/63/044/002/009/065
B102/B186

the γ -rays were detected by a NaI(Tl) crystal ($2 \cdot 10^{-7}$ sec) connected with an ФЭУ-29 (FEU-29) photomultiplier; the neutron detector ($4 \cdot 10^{-9}$ sec) was provided with an FEU-24. The coincidence between 4.43-Mev γ -quanta ($2^+ \rightarrow 0^+$ transition) and the neutrons scattered through a certain angle was recorded by scintillation counters. The target was a graphite cylinder 15 cm high and 6.5 cm thick. The γ -ray angular distributions were measured in the n - n' plane for the fixed n' emission angles 40 and 135°. They can be described by $f(\theta_\gamma) = 1 + b \sin^2(\theta_\gamma - \theta_0)$ where b and θ_0

were calculated by the method of least squares. The scattering cross-section ratio was $\sigma(40^\circ)/\sigma(135^\circ) = 1.8 \pm 0.4$. When the results obtained for neutrons are compared with the analogous ones for protons it can be seen that for small angles the scattering mechanism depends only slightly on the nature of the nucleon, but for large angles the results obtained for neutrons differ from those for protons. It cannot yet be decided if the direct scattering mechanism is always contributed by another mechanism nor to what extent this occurs. For large scattering angles this contribution will be very important. There are 6 figures and 1 table.

ASSOCIATION: Fizicheskii institut im. P. N. Lebedeva Akademii nauk SSSR
(Institute of Physics imeni P. N. Lebedev of the Academy of Sciences USSR)

Card 2/13

GOL'DANSKIY, V.I.; KUZNETSOV, B.G., prof.; MIGDAL, A.B.; FRANK, I.M.; CHERNOV, A.G.; FAYNBOY, I.B., red.

[The constitution of matter; first talk] Stroenie veshchestva; beseda pervaya. [By] V.I. Gol'danskii i dr. Moskva, Izd-vo "Znanie," 1964. 38 p. (Novoe v zhizni, nauke, tekhnike. IX Seriya: Fizika, matematika, astronomia, no.5) (MIRA 17:5)

1. Chleny-korrespondenty AN SSSR (for Gol'danskiy, Migdal, Frank).

FRANK, I. M.; BUNIN, B. N.; NIKOLAYEV, S. K.; SHABALIN, Ye. P.; SHAPIRO, F. L.

"The experience of the pulsed fast reactor operation and its characteristics at injection of neutrons from a microtron."

report submitted for 3rd Intl Conf, Peaceful Uses of Atomic Energy, Geneva, 31 Aug-9 Sep 64.

ACCESSION NR: AT4041826

S/2504/64/024/000/0203/0211

AUTHOR: Frank, I. M.

TITLE: Some characteristics of the elastic moderation of neutrons

SOURCE: AN SSSR. Fizicheskij Institut. Trudy*, v. 24, 1964. Issledovaniya po neytronnoy fizike (Research in neutron physics), 203-211

TOPIC TAGS: neutron moderation, neutron, neutron velocity loss, age diffusion theory, dispersion spectrum, age equation, elastic moderation, neutron spectrum

ABSTRACT: In the investigation of the moderation and diffusion of neutrons, it is important to solve the problem of the instantaneous neutron spectrum. The age-diffusion theory assumes that neutron energy loss in scattering on heavy nuclei is small and that the neutron energy is a continuous function of time. However, there are deviations in energy and velocity from the mean square values which one would expect, for example, if all neutrons had the same initial energy. These deviations in the instantaneous neutron spectrum are usually not considered because of the complexity of the non-stationary theory.

Card 1/5

ACCESSION NR: AT4041826

However, on the basis of a simple and approximate approach, the square of the velocity deviation from the mean value can be calculated. Starting from the probability equation that a neutron of velocity V (energy E) after a collision will have a velocity in the interval from V' to $V' + dV'$ (energy E' to $E' + dE'$), the average velocity so obtained is given by

$$\bar{v} = \int_{v_{\min}}^v v' W(v') dv' = \frac{M + \frac{1}{3M}}{M + 1} v \quad (1)$$

The velocity loss as a function of time is given by

$$\frac{dv}{dt} = -\frac{1 - \frac{1}{3M}}{M + 1} \frac{v}{\lambda} = -\frac{1}{M\lambda} v^2 \quad (2)$$

where M' is the mass of the nucleus. The mean square of the neutron velocity after one collision is then given by

$$\overline{v^2} = \int_{v_{\min}}^v v'^2 W(v') dv' = \frac{M^2 + 1}{(M + 1)^2} v^2 \quad (3)$$

Card 2/5

ACCESSION NR: AT4041826

and the square of the mean square deviation in velocity Δv^2 is given by

$$[\Delta v^2]_{\text{mean}} = (\overline{v'^2} - \overline{v'}^2) = \overline{v'^2} - (\overline{v'})^2 = \frac{1 - \frac{1}{3M^2}}{3(M+1)^2} v^2. \quad (4)$$

The time-dependent changes in values of $(\frac{\Delta v^2}{v^2})$ due to the fluctuation in the mean free path is given by

$$\frac{d}{dt} \left(\frac{\Delta v^2}{v^2} \right)_{\text{cr}} = \frac{d}{dt} \frac{\Delta v_1^2}{v^2} + \frac{d}{dt} \frac{\Delta v_2^2}{v^2} = \frac{4}{3M^2} \frac{v}{\lambda}. \quad (5)$$

If the velocity loss were always equal to a mean value, the change in $\frac{\Delta v^2}{v^2}$ would occur according to

$$\frac{d}{dt} \left(\frac{\Delta v^2}{v^2} \right)_{\text{cr}} \approx -\frac{2v}{M\lambda} \frac{\Delta v^2}{v^2}. \quad (6)$$

It is shown here that the instantaneous neutron spectrum converges toward the stationary distribution given by

$$\frac{\Delta v^2}{v^2} = \frac{2}{3M^2}. \quad (7)$$

Card 3/5

ACCESSION NR: AT4041826

The interval X^2 of the change in velocity needed to establish a stationary value of $\frac{\Delta v^2}{v^2}$ is calculated and expressed as

$$x^2 = \left(\frac{v}{v_0}\right)^2 \left(x_0^2 - \frac{2}{3M'}\right) + \frac{2}{3M'} \quad (8)$$

If the moderator contains an impurity with lighter nuclei with effective mass $\frac{M'}{2}$, the time change in velocity is given by

$$\left(\frac{\Delta v^2}{v^2}\right) \approx \frac{2}{3M'} \left(1 + \frac{M'\lambda}{M_1\lambda_1}\right) \quad (9)$$

and that in the magnitude $\frac{\Delta v^2}{v^2}$ is given by

$$\frac{dv}{dt} = - \left(\frac{1}{M'} \frac{v^2}{\lambda} + \frac{1}{M_1} \frac{v^2}{\lambda_1} \right) = - \frac{v^2}{M'\lambda} \left(1 + \frac{M'\lambda}{M_1\lambda_1} \right) \quad (10)$$

4/5

Card

ACCESSION NR: AT4041826

Finally, the conditions for the validity of the age-diffusion equation are discussed.
"The author is indebted to F. L. Shapiro and M. V. Kazarnovkiy for numerous valuable
comments." Orig. art. has: 40 formulas.

ASSOCIATION: Fizicheskiy institut im. P. N. Lebedeva, AN SSSR (Physics Institute,
AN SSSR)

SUBMITTED: 00

ENCL:00

SUB CODE: NP

NO REF SOV: 004

OTHER: 000

Card 5/5

L 5259-66 EWT(1)/EWT(m)/EPF(c)/ETC/EPF(n)-2/EWG(m) TJP(c) WW
 ACC NR: AP5014805 SOURCE CODE: UR/0030/65/000/005/0396/0098

AUTHOR: Frank, I. M. (Corresponding member AN SSSR) 5/

ORG: none 14
B

TITLE: Scientific session on the application of nuclear methods to solid-state physics

SOURCE: AN SSSR. Vestnik, no. 5, 1965, 96-98 2/14

TOPIC TAGS: nuclear physics conference, solid state physics conference, solid state physics, nuclear physics, nonmilitary nuclear application

ABSTRACT: A meeting of nuclear scientists and scientists from other fields was organized by the Department of General and Applied Physics and the Department of Nuclear Physics (both of the Academy of Sciences USSR) and the State Committee on the Use of Atomic Energy USSR and held from 9-10 December 1964 in Moscow. 44.55

The program consisted of reports on problems common to the physics of solids and liquids and nuclear physics. Yu. M. Kagan reported on the Mossbauer effect as a means for detecting very slight differences in nuclear frequencies and studying the internal fields of atoms or internal motions of matter. F. L. Shapiro discussed various uses of slow neutron 44.55

Card 1/2

L 5259-66

ACC NR: AP5014805

scattering and described nuclear reactors¹⁹ as important devices for the study of condensed media. Positron annihilation as a means for investigating the physical and chemical properties of substances was treated in a report by V. I. Gol'danskiy. B. N. Samoylov's report discussed the loss of parity in weak interactions as a means for studying fields within matter with polarized nuclei. I. M. Frank reported on the possibilities inherent in the interdependence of transition radiation and the optical properties of the substance.

The necessity for closer cooperation between nuclear physicists and solid-state specialists is called for and the emergence of a new discipline—nuclear solid-state physics—is foreseen. The session, which was limited to three meetings, did not cover such problems as nuclear paramagnetism and radiation physics. These topics will be studied in future sessions. [FSB: v. 1, no. 12]

SUB CODE: NP, SS / SUBM DATE: none

CC
Card 2/2

FRANK, I.M.

Transient radiation and the optical properties of matter.
Usp. fiz. nauk 87 no.2:189-210 0 '65. (MIRA 18:11)

S/276/63/000/002/006/052
A052/A126

AUTHORS: Frank, János, and Kincses, István

TITLE: A method of manufacturing steel and non-ferrous balls

PERIODICAL: Referativnyy zhurnal, Tekhnologiya mashinostroyeniya, no. 2, 1963, 39, abstract 2B152 P. (Hung. pat., cl. 49 1, no. 148840, December 31, 1961)

TEXT: The antifriction bearing plant in Debrecen (Hungary) has patented a method of manufacturing balls. The essence of the method is that the wire is heated in the voltaic arc, in an acetylene flame, by the electric resistance method or induction, to the temperature over the melting point and sprayed by compressed air. Fused drops fall into a corresponding cooling medium (oil, kerosene, aqueous solution of some salt or emulsion) and owing to the surface tension solidify in the form of balls. The process makes it possible to produce hardened balls of various structures by a proper selection of the cooling medium. The described process can be realized by means of the spraying pistol.

(Abstracter's note: Complete translation.) V. Bachin

Card 1/1

FRANK, Janos; DREDAN, Istvan

Who should be in charge of manufacturing grinding wheels? Musz
elet 19 no.8:4 9 Ap '64.

FRANK, Jaroslav

The Roirant R7 machine.

p. 302 (Sklar A Keramik) Vol. 7, no. 10, Oct. 1957, Fraha, Czechoslovakia

SO: MONTHLY INDEX OF EAST EUROPEAN ACCESSIONS (EEAI) LC, VOL. 7, NO. 1, JAN. 1958

FRANK, JAROSLAV

CZECHOSLOVAKIA/Chemical Technology - Chemical Products and
Their Application - Ceramics, Glass, Binders,
Concrete.

H-13

Abs Jour : Ref Zhur - Khimiya, No 3, 1958, 8756

Author : Frank Jaroslav, Obraz Karel

Inst : -

Title : Container Glass for Canning Factories.

Orig Pub : Sklar a keramik, 1957, 7, No 4, 110-112

Abstract : In manufacturing modern container glass it is necessary to take into account the technological process of canning and the stresses to which are subjected the glass jars in the course thereof. The jars are subjected to the greatest stresses during the process of sterilization. The conditions of annealing of the glass must ensure the elimination of internal stresses above $95 \text{ m} \mu / \text{cm}$. Problems are considered which relate to design of jar necks, types of covers and color of glass depending on the nature of the preserved products.

Card 1/1

FRANK, J.

Preparing the standardization of devices for closing bunkers. p. 81.

Vol. 3, no. 3, Mar. 1954 (Mechanisace)
INZENYRSKE STAVBY
Praha, Czechoslovakia

So: Eastern European Accession Vol. 5 No. 4 April 1956

FRANK, J.

Use of a claw-type loader in building. p. 127.

Vol. 3, no. 4, April 1954 (Mechanisation)
INZENYRSKE STAVBY
Praha, Czechoslovakia

So: Eastern European Accession Vol. 5 No. 4 April 1956